Sobotta

Atlas of Human Anatomy

Edited by R. Putz and R. Pabst
In collaboration with Renate Putz

Translation by S. Bedoui

Volume 2
Trunk, Viscera, Lower Limb

14th edition, newly edited
704 colour plates with 914 figures



URBAN & FISCHER
München · Jena

Sobotta

Atlas of Human Anatomy

Volume 2
Trunk, Viscera, Lower Limb

Correspondence and feedback should be addressed to:
Elsevier GmbH, Urban & Fischer Verlag, Department for Medical Student Information, Alexander Gattnarzik, Karlstraße 45, 80333 Munich, Germany
e-mail: medizinstudium@elsevier.de

Addresses of the editors:

Professor Dr. med. Reinhard Putz
Vorstand des Anatomischen Instituts
der Ludwig-Maximilians-Universität
Pettenkoferstraße 11
80336 München
Germany
e-mail: reinhard.putz@med.uni-muenchen.de

Professor Dr. med. Reinhard Pabst
Leiter der Abteilung
Funktionelle und Angewandte Anatomie
Medizinische Hochschule Hannover
Carl-Neuberg-Straße 1
30625 Hannover
Germany
e-mail: pabst.reinhard@mh-hannover.de

Bibliographic information published by Die Deutsche Bibliothek

Die Deutsche Bibliothek lists this publication in the Deutsche National-bibliografie; detailed bibliographic data is available in the Internet at http://dnb.ddb.de.

All rights reserved

14th edition 2006
©2006 Elsevier GmbH, Munich
Urban & Fischer Verlag is an imprint of Elsevier GmbH.

05 06 07 08 09 5 4 3 2 1

All rights, including translation, are reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any other form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of the publisher.

The 14th edition of the Sobotta Atlas consists of two volumes and a booklet containing tables (enclosed in Volume 1):

Volume 1: Head, Neck, Upper Limb

Volume 2: Trunk, Viscera, Lower Limb

Translation: Dr. med. Sammy Bedoui, Hannover
Editorial staff at Elsevier: Dr. med. Dorothea Hennessen
Alexander Gattnarzik
Dr. rer. nat. Andrea Richarz
Illustrators: Ulrike Brugger, Munich; Rüdiger Himmelhan, Heidelberg; Horst Ruß, Munich; Henriette Rintelen, Velbern
Book production: Renate Hausdorf, Munich
Composed by: Typodata, Munich
Printed and bound by: Appl, Wemding
Cover design: Carsten Tschirner, Munich
Printed on Nopacoat 115 g

Printed in Germany
ISBN-13: 978-0-443-10349-0
ISBN-10: 0-443-10349-6

This atlas was founded by Johannes Sobotta†, former Professor of Anatomy and Director of the Anatomical Institute of the University of Bonn, Germany.

German Editions:
1st Edition: 1904–1907 J. F. Lehmanns Verlag, Munich
2nd–11th Edition: 1913–1944 J. F. Lehmanns Verlag, Munich
12th–20th Edition: 1948–1993 Urban & Schwarzenberg, Munich
13th Edition: 1953, editor H. Becher
14th Edition: 1956, editor H. Becher
15th Edition: 1957, editor H. Becher
16th Edition: 1967, editor H. Becher
17th Edition: 1972, editors H. Ferner and J. Staubesand
18th Edition: 1982, editors H. Ferner and J. Staubesand
19th Edition: 1988, editor J. Staubesand
20th Edition: 1993, editors R. Putz and R. Pabst
21st Edition: 2000, editors R. Putz and R. Pabst, Urban & Fischer Verlag, Munich
22nd Edition: 2006, editors R. Putz and R. Pabst, Elsevier GmbH, Munich

Foreign Editions:
Arabic Edition
Modern Technical Center, Damascus
Chinese Edition (complex characters)
Ho-Chi Book Publishing Co, Taiwan
Chinese Edition (simplified Chinese edition)
Elsevier, Health Sciences Asia, Singapore
Croatian Edition
Naklada Slap, Jastrebarsko
Dutch Edition
Bohn Stafleu van Loghum, Houten
English Edition (with nomenclature in English)
Atlas of Human Anatomy
Lippincott Williams & Wilkins
English Edition (with nomenclature in Latin)
Atlas of Human Anatomy
Elsevier GmbH, Urban & Fischer
French Edition
Atlas d'Anatomie Humaine
Tec & Doc Lavoisier, Paris
Greek Edition (with nomenclature in Greek)
Maria G. Parissianos, Athens
Greek Edition (with nomenclature in Latin)
Maria G. Parissianos, Athens
Hungarian Edition
az ember anatómiájának atlasza
Alliter Kiadái, Budapest
Indonesian Edition
Atlas Anatomi Manusia
Penerbit Buku Kedokteran EGC, Jakarta
Italian Edition
Atlante di Anatomia Umana
UTET, Torino
Japanese Edition
Igaku Shoin Ltd., Tokyo
Korean Edition
ShingHeung MedScience, Seoul
Polish Edition
Atlas anatomii cztowieka
Urban & Partner, Wroclaw
Portuguese Edition (with nomenclature in English)
Atlas de Anatomia Humana
Editora Guanabara Koogan, Rio de Janeiro
Portuguese Edition (with nomenclature in Latin)
Atlas de Anatomia Humana
Editora Guanabara Koogan, Rio de Janeiro
Spanish Edition
Atlas de Anatomia Humana
Editorial Medica Panamericana, Buenos Aires/Madrid
Turkish Edition
Insan Anatomisi Atlasi
Beta Basim Yayim Dagitim, Istanbul

Contents

Preface VI

Back 1–45

Surface anatomy 1
Vertebral column, Columna vertebralis 2–25
Muscles of the back, Musculi dorsi, suboccipital muscles, Musculi suboccipitales, and muscles of the neck, Musculi colli 26–34
Computed tomographic sections through the vertebral column 35
Vessels and nerves of the back and the occipital region 36–40
Vessels and nerves of the vertebral canal 41–45

Thoracic and abdominal wall 46–73

Surface anatomy 46
Ribs, Costae, thoracic cavity, Cavea thoracis 47–49
Sternum, Sternum 50–51
Muscles of the thorax, Musculi thoracis, and abdominal muscles, Musculi abdominis 52–59
Inguinal canal 60–61
Diaphragm, Diaphragma 62–63
Breast, Mamma 64–65
Vessels and nerves of the thoracic and abdominal wall 66
Segmental sensory innervation 67
Lumbosacral plexus, Plexus lumbosacralis 68
Anterior abdominal wall 69–71
Sections through the abdominal wall 72–73

Thorax 74–127

Heart, Cor 74–89
Trachea, Trachea 90–91
Lungs, Pulmones 92–100
Shape of the heart 101
Bronchoscopy 102
Projection of trachea, lungs and pleural borders 103
Oesophagus, Oesophagus 104–109
Thymus, Thymus 110–111
Pleural cavity, Cavitas pleuralis, and mediastinum, Mediastinum 112–113
Aorta, Aorta 114
Posterior mediastinum 115–116
Autonomous nervous system, Pars autonomica 117
Phrenic nerve, Nervus phrenicus 118
Sections through the thorax 119–127

Abdominal viscera 128–181

Stomach, Gaster 128–133
Small intestine, Intestinum tenue 134–137
Large intestine, Intestinum crassum 138–141
Liver, Hepar, gall bladder, Vesica biliaris, and bile duct system 142–151
Pancreas, Pancreas 152–155
Spleen, Splen [Lien] 156
Peritoneal cavity, Cavitas peritonealis, development .. 157
Location of the abdominal viscera, Situs viscerum 158–167
Abdominal arteries, overview 168
Abdominal veins, overview 169
Coeliac trunc and superior as well as inferior mesenteric arteries and veins, Truncus coeliacus, Arteriae et Venae mesentericae superior et inferior 170–178
Vessels and nerves of the retroperitoneal space 179
Hepative portal vein, Vena portae hepatis 180–181

Pelvic viscera and retroperitoneal space 182–261

Kidney, Ren [Nephros] 182–188
Suprarenal gland, Glandula suprarenalis 189
Urinary bladder, Vesica urinaria 190–191
Male urinary and genital organs 192–199
Female urinary and genital organs 200–209
Rectum, Rectum 210–215
Renal artery, Arteria renalis 216
Organs of the retroperitoneal space 217
Vessels and nerves of the retroperitoneal space 218–221
Innervation of male genitalia 222
Innervation of female genitalia 223
Internal pelvic arteries 224–225
Pelvic viscera 226–229
Pelvic diaphragm and perineum, male and female external genitalia 230–241
Sections through abdomen and pelvis 242–261

Lower limb 262–384

Surface anatomy 262
Bones, joints, ligaments 263–305
Muscles of the lower limb 306–343
Vessels and nerves 344–379
Sections through the lower limb 380–384

Appendix 385–399

Index 385–399

Preface

It was just over a hundred years ago that Johannes Sobotta set out to publish the first edition of his Atlas of Human Anatomy. Since then, this piece of work has evolved step by step as a result of the constant interaction between students, lecturers and editors. It has not only been the most modern basis for the complex subject of macroscopic anatomy throughout many generations of doctors, but has also developed into a lasting work of reference for both clinical training and advanced medical education. All in all, it has become a book for a medical doctor's life. Once again, in this new edition the additional figures have been drawn strictly on the basis of original specimens.

The 14th edition has been particularly designed to meet the demands of a reformed medical curriculum, emphasizing the integration of clinical medicine into the preclinical curriculum. For this purpose, the new edition has been extended to include the following features:

- Surface anatomy including projection of internal organs (45 colour photos)
- Anatomical diagrams next to imaging figures
- Integration of imaging techniques to a greater extent (ultrasound, X-ray, CT, MRI; 119 figures)
- Endoscopic, intraoperative colour images and figures exemplifying techniques of puncture and examination (54 figures)
- Images of patients presenting with typical palsies
- Diagrams of the most important arterial variations (93 figures)
- Frequent variations in the location of internal organs (24 figures)
- Integration of histology at low magnification of important internal organs (intestine, liver, kidney, etc.)

In order to improve the presentation of the knowledge, the following features have been introduced:

- Clear-cut arrangement of the chapters according to the different regions of the body
- Thematically corresponding figures presented on double pages
- A concise, separate booklet (included in Volume 1) contains tables of muscles, joints and nerves, enabling the reader to place it next to any figure in the atlas

One particular aim of the new edition is to facilitate finding of specific structures. The SOBOTTA depicts anatomical structures precisely without the reader loosing the greater picture. Therefore, specific didactic tools have been improved and new aspects included:

- Each chapter has been allocated to a particular colour
- A "menu bar" on each double page ensures precise orientation within a given chapter
- The number of outlines depicting spatial orientation has been significantly increased (270 figures)
- Overviews of total body regions ensure general orientation
- New diagrams of particular muscles clarify their location and course (24 figures)
- Confusion is kept to a minimum by only depicting limbs of the right side of the body
- "Compass roses" point to adjacent figures, thus facilitating following a given structure over several pages
- Continuous leader lines facilitate finding of structures
- Coloured dots at the end of leader lines in topographic diagrams mark arteries, veins, nerves, and muscles
- The figures in the booklet relate directly to the figures in the atlas
- The larger dimensions of the book improve clarity

With the exception of discussions about the general concept of the atlas and mutual correction, the editors have worked separately on individual chapters, with the work divided as follows:

R. Putz: General anatomy, upper limb, brain, eye, ear, back, lower limb;
R. Pabst: Head, neck, thoracic and abdominal walls, thoracic, abdominal and pelvic viscera.

The inclusion of a large number of new figures is the result of the extraordinary capability of the following medical illustrators: Ulrike Brugger, Rüdiger Himmelhan and Horst Ruß. It is to their credit that the classic "SOBOTTA style" has been retained. Several of the diagrams have been generated on the computer by Henriette Rintelen. We also gratefully acknowledge our clinical colleagues for making clinical illustrations available to us (see picture credits). We owe a debt of gratitude to our colleagues from the institutes for their understanding and helpful suggestions. Dr. N. Sokolov and A. Buchhorn have put meticulous efforts into generating the specimen preparations. S. Fryk and G. Hoppmann have supported us in text processing.

The staff of the editorial office of Elsevier publishers, in particular Dr. D. Hennessen and A. Gattnarzik, has our sincere thanks. Some of the creative development of the work is a result of very fruitful discussions. We would also like to thank R. Hausdorf for tremendous efforts in the production of the atlas. R. Putz together with G. Meier were responsible for the proofreading and simplification of page design and legends. Our special thanks go to Dr. U. Osterkamp-Baust for generating the index, and all others involved in the corrections. With our joint efforts, the SOBOTTA has been once more modernized both in contents and design.

We have included many of the helpful suggestions made over the years by students and colleagues, and would therefore ask all readers of this edition to pass on to us any criticism or suggestions on the new format of this atlas.

Munich and Hannover, September 2005
R. Putz and R. Pabst

Univ.-Prof. Dr. med. Reinhard Putz

Born in Innsbruck/Austria

1962–1968	Studied medicine at the Leopold-Franzens-University of Innsbruck
1968	Received a doctorate
1968–1982	University assistant at the Institute of Anatomy at the University of Innsbruck
1978	Lecturer in anatomy
1979	Consultant for anatomy
1982–1989	Chair of the Anatomical Institute at the Albert-Ludwigs-University of Freiburg
since 1989	Chair of the Anatomical Institute at the Ludwig-Maximilians-University of Munich
1992–1994	President of the European Association of Clinical Anatomists
1993	Registration to practise medicine
1998–1999	Chairman of the Anatomical Society
1999	Member of the Akademie der Naturforscher und Ärzte (Leopoldina)
2002	Dr. h.c. of the University of Constanta, Romania
2003	Prorector I of the Ludwig-Maximilians-University of Munich

Research and fields of interest

- Functional anatomy of the passive locomotor system
- Evolution and functional anatomy of the vertebral column
- Form-function-relations of joints
- Applied anatomy (anatomical basics of orthopaedics, surgery, radiology)
- Questions about the contents and organisation of the medical curriculum
- Development of didactic training programmes at universities

Univ.-Prof. Dr. med. Reinhard Pabst

Born in Posen, grown up in Lüneburg

1965–1970	Studied medicine at the Hannover Medical School, and in Glasgow/Scotland
1970	University degree and doctorate
1971	Registration to practise medicine
1971–76	Scientific associate in the Department of Clinical Physiology, University of Ulm
1976	Lecturer for Clinical Physiology, University of Ulm, and new lectureship at the Hannover Medical School
1976–80	Senior assistant in the Department of Functional and Applied Anatomy, Hannover Medical School
1978	Extension of the Venia legendi to include anatomy
1980–1992	Head of the Department of Topographic Anatomy and Biomechanics
since 1992	Head of the Department of Functional and Applied Anatomy, Hannover Medical School
1986–1990	Prorector for studies and education, Hannover Medical School
1993–1997	Dean of the Hannover Medical School
1997–1998	Chairman of the Anatomical Society
1999–2003	Prorector of Research of the Hannover Medical School
2001	Member of the Akademie der Naturforscher und Ärzte (Leopoldina)

Research and fields of interest

- Functional anatomy of lymphatic organs
- Proliferation and migration of lymphocytes
- Development of the intestinal immune system
- Function of the pulmonary immune system
- Questions of a clinically orientated anatomy in the medical curriculum
- Evaluation of teaching

Picture credits

The editors sincerely thank all clinical colleagues that made ultrasound, computed tomographic and magnetic resonance images as well as endoscopic and intraoperative pictures available:

Prof. Altaras, Centre for Radiology, University of Gießen (Fig. 1009, 1031, 1032)
Dr. Baumeister, Department of Radiology, University of Freiburg (Fig. 1157)
PD Dr. Burgkardt, Orthopaedic Clinic, Technical University of Munich (Fig. 1431)
Prof. Brückmann & Dr. Linn, Neuroradiology, Institute for Diagnostic Radiology, University of Munich (Fig. 662, 726 a, b, 727 a, b)
Prof. Daniel, Department of Cardiology, University of Erlangen (Fig. 899, 900, 901, 973)
Prof. Degenhardt, Bielefeld (Fig. 1133, 1135)
Prof. Galanski & Dr. Kirchhoff, Department of Diagnostic Radiology, Hannover Medical School (Fig. 962, 964, 1212, 1213)
Prof. Galanski & Dr. Schäfer, Department of Diagnostic Radiology, Hannover Medical School (Fig. 871a, b, 927, 971, 991, 1194, 1204, 1207, 1209)
Prof. Gebel, Department of Gastroenterology, Hepatology and Endocrinology, Hannover Medical School (Fig. 242, 1011, 1026, 1027, 1038, 1039, 1074, 1092)
Dr. Goei, Radiology, Heerlen, The Netherlands (Fig. 1150, 1151) (with permission from Radiology 173; 137–141: 1989)
Dr. Greeven, St.-Elizabeth-Hospital, Neuwied (Fig. 150, 1238)
Prof. Hoffmann & PD Dr. Bektas, Clinic for Abdominal and Transplantation Surgery, Hannover Medical School (Fig. 1028, 1040)
Prof. Hohlfeld, Clinic for Pneumology, Hannover Medical School (Fig. 929, 930)
Prof. Jonas, Urology, Hannover Medical School (Fig. 1102 a, b, 1103)
Prof. Kampik & Prof. Müller, Ophthalmology, University of Munich (Fig. 657)
Dr. Kirchhoff & Dr. Weidemann, Department of Diagnostic Radiology, Hannover Medical School (Fig. 1041, 1093, 1102, 1196, 1198, 1200, 1202, 1205)
Prof. Kremers, Department for Restorative Dentistry and Periodontology, University of Munich (Fig. 169)
Prof. Kunze, von Haunersches Children's Hospital, University of Munich (Fig. 327–330)
Dr. Meyer, Department of Gastroenterology, Hepatology and Endocrinology, Hannover Medical School (Fig. 944, 987 a, b, 992, 1145, 1146)
Prof. Müller-Vahl, Neurology, Hannover Medical School (Fig. 128 a, b)
Prof. Pfeifer, Radiology, Institute for Diagnostic Radiology, University of Munich (Fig. 293, 294, 310, 312, 449, 451, 769–772, 807–810, 1255, 1286, 1287, 1317, 1318)
PD Dr. Rau, Department of Radiology, University of Freiburg (Fig. 912, 925, 926)
Prof. Ravelli †, formerly Institute of Anatomy, University of Innsbruck (Fig. 767)
PD Dr. Rieger, Radiology, Institute for Diagnostic Radiology, University of Munich (Fig. 1395)
Prof. Reich, Orofacial Surgery, University of Bonn (Fig. 113 a, b)
Prof. Reiser & Dr. Wagner, Institute for Diagnostic Radiology, University of Munich (Fig. 436, 449, 451, 453, 571, 572, 573, 576, 577, 792)
Prof. Rudzki-Janson, Department of Orthodontics, University of Munich (Fig. 72, 73)
Dr. Scheibe, Department of Surgery, Rosman Hospital, Breisach (Fig. 1289 a–c)
Prof. Scheumann, Clinic for Abdominal and Transplantation Surgery, Hannover Medical School (Fig. 243, 244, 245)
Prof. Schillinger, Department of Gynaecology, University of Freiburg (Fig. 1134)
Prof. Schliephake, Orofacial Surgery, Göttingen (Fig. 152, 196, 197)
Prof. Schlösser, Centre for Gynaecology, Hannover Medical School (Fig. 1132a, b, 1140, 1141, 1178)
Prof. Schumacher, Neuroradiology, Department of Radiology, University of Freiburg (Fig. 461 a, b)
Dr. Sommer & PD Dr. Bauer, Radiologists, Munich (Fig. 662, 1290–1292)
Prof. Stotz, Paediatrics, University of Munich (Fig. 1239, 1240)
PD Dr. Vogl, Radiology, University of Munich (Fig. 436, 453, 644, 645)
Prof. Vollrath, Ear-Nose-Throat Department, Mönchengladbach (Fig. 229, 230, 231)
Prof. Wagner †, Diagnostic Radiology II, Hannover Medical School (Fig. 952, 1064, 1065, 1069, 1072, 1152)
Prof. Wenz, formerly Department of Radiology, University of Freiburg (Fig. 768)
Prof. Witt, Department of Neurosurgery, University of Munich (Fig. 405)
Dr. Willführ, formerly Clinic for Abdominal and Transplantation Surgery, Hannover Medical School (Fig. 1051)
PD Dr. Wimmer, Department of Radiology, University of Freiburg (Fig. 799)

Additional illustrations were obtained from the following textbooks:

Benninghoff-Drenckhahn: Anatomie, Band 1 (Drenckhahn D., Hrsg.), 16. Aufl., Urban & Fischer, München 2003 (Fig. 823, 824, 848, 849, 850)
Welsch, U.: Lehrbuch Histologie, Urban & Fischer, München 2003 (Fig. 922, 1020, 1097, 1122, 1123)
Welsch, U. (Hrsg.): Sobotta, Atlas Histologie, 6. Aufl., Urban & Fischer, München 2002 (Fig. 1037, 1085, 1125 a, b)
Wicke, L.: Atlas der Röntgenanatomie, 3. Aufl., Urban & Schwarzenberg, München–Wien–Baltimore 1985 (Fig. 943 a)



Fig. 728 Back, Dorsum; surface anatomy.



Fig. 729 Regions and orientation lines of the back.

Skeleton of the trunk



Fig. 730 Skeletal system, Systema skeletale, of the trunk.



Fig. 731 Vertebral column, Columna vertebralis; pectoral girdle, Cingulum pectorale, and pelvic girdle, Cingulum pelvicum; median section through the vertebral column; medial view.



Fig. 732 Vertebral column, Columna vertebralis; pectoral girdle, Cingulum pectorale, and pelvic girdle, Cingulum pelvicum; median section through the vertebral column; viewed from the left.

Vertebral column



Fig. 733 Vertebral column, Columna vertebralis; ventral view.

Fig. 734 Vertebral column, Columna vertebralis; dorsal view.



Fig. 735 Vertebral column, Columna vertebralis; viewed from the left.





Fig. 736 a–e Regional characteristics of the vertebrae. Only in the thoracic region of the vertebral column do the lateral parts (labelled in red) remain separated and form ribs.

a First cervical vertebra, Atlas
b Fourth cervical vertebra, Vertebra cervicalis IV
c First thoracic vertebra, Vertebra thoracica I, with corresponding ribs, Costae, and sternum, Sternum
d Third lumbar vertebra, Vertebra lumbalis III
e Sacrum, Os sacrum



Fig. 737 Vertebral development.
Demonstration of the appearance of primary ossification centres in a lumbar vertebra (pedicle: second foetal month; corpus: third to sixth foetal month).
Synostosis of the ossification centres of the vertebral arch with those of the corpus occurs between the third and the sixth year of life.



Fig. 738 Vertebral development.
Circular ossification centres (= rims*) appear in the epiphyses of the vertebrae during the eighth year of life and fuse with the vertebral bodies until the eighteenth year of life.
The central portions of the epiphyses remain as hyaline cartilaginous laminae ** throughout life.
Secondary ossification centres (apophyses) develop at the processes.

Atlas and axis



Fig. 739 Occipital bone, Os occipitale; section illustrating the foramen magnum and the articular surfaces of the atlanto-occipital joint; inferior view.



Fig. 740 First cervical vertebra, Atlas; superior view.
The superior articular surfaces of the atlas are frequently divided.
* Canalis arteriae vertebralis as a variation



Fig. 741 First cervical vertebra, Atlas; inferior view.



Fig. 742 First and second cervical vertebrae, Atlas et Axis; median section.



Fig. 743 Second cervical vertebra, Axis; ventral view.



Fig. 744 Second cervical vertebra, Axis; dorsosuperior view.



Fig. 745 Fifth cervical vertebra, Vertebra cervicalis V; superior view.
The tips of the spinous processes of the cervical vertebrae II–VI are frequently divided.



Fig. 746 Seventh cervical vertebra, Vertebra cervicalis VII; superior view.
In general the seventh cervical vertebra can be easily identified by its protruding spinous process and is referred to as vertebra prominens. In fact, the spinous process of the first thoracic vertebra often protrudes even further.



Fig. 747 Second to seventh cervical vertebrae, Vertebrae cervicales II–VII; ventral view.



Fig. 748 First to seventh cervical vertebrae, Vertebrae cervicales I–VII; dorsolateral view.

Thoracic and lumbar vertebrae



Fig. 749 Vertebra, Vertebra; typical structural features exemplified by the fifth thoracic vertebra; superior view.
* Also: rim of vertebral body



Fig. 750 Sixth thoracic vertebra, Vertebra thoracica VI; viewed from the left.



Fig. 751 Tenth thoracic vertebra, Vertebra thoracica X; ventral view.



Fig. 752 Twelfth thoracic vertebra, Vertebra thoracica XII; viewed from the left.
* Region of the vertebral arch between the superior and inferior articular processes ("isthmus" = interarticular portion)



Fig. 753 Third lumbar vertebra, Vertebra lumbalis III; median section; specimen of an older individual.
* Ossification of the ligamentous insertions



Fig. 754 Tenth to twelfth thoracic vertebrae, Vertebrae thoracicae X–XII, and first to second lumbar vertebrae, Vertebrae lumbales I–II; dorsal view from the left.



Fig. 755 Fourth lumbar vertebra, Vertebra lumbalis IV; superior view.



Fig. 756 Fourth lumbar vertebra, Vertebra lumbalis IV; ventral view.



Fig. 757 Fifth lumbar vertebra, Vertebra lumbalis V; median section.
Note the characteristic wedge-shape of the body of the fifth lumbar vertebra.

* Region of the vertebral arch between the superior and inferior articular process. Here in the fifth and less frequently in the fourth lumbar vertebra a cleft, bridged by connective tissue (spondylolysis) can be formed. This is probably caused by local bending stress. As a consequence, the superior vertebra may slip (olisthesis) onto the inferior vertebra (spondylolisthesis).

Sacrum



Fig. 758 Sacrum, Os sacrum; dorsal view.



Fig. 759 Sacrum, Os sacrum; ventroinferior view.



Fig. 760 Sacrum, Os sacrum; the bone has been sectioned at the level of the second sacral vertebra; superior view.



Fig. 761 Sacrum, Os sacrum; viewed from the right.



Fig. 762 Sacrum, Os sacrum; median section.
* Remnants of intervertebral disc tissue persist even in the adult.



Fig. 763 Sacrum, Os sacrum; gender differences.



Fig. 764 Sacrum, Os sacrum; gender differences.



Fig. 765 Coccyx, Os coccygis; ventrosuperior view.



Fig. 766 Coccyx, Os coccygis; dorsoinferior view.

Cervical part of the vertebral column, radiography



Fig. 767 Cervical vertebrae, Vertebrae cervicales; lateral radiograph of the cervical part of the vertebral column; upright position; the central beam is directed onto the third cervical vertebra; shoulders are pulled downwards.



Fig. 768 Cervical vertebrae, Vertebrae cervicales; AP-radiograph of the cervical part of the vertebral column; upright position; the central beam is directed onto the third cervical vertebra.
* Intervertebral disc spaces

Thoracic part of the vertebral column, radiography



Fig. 769 Thoracic vertebrae, Vertebrae thoracicae; lateral radiograph of the thoracic part of the vertebral column; upright position with the thorax in inspiration; the central beam is directed onto the sixth thoracic vertebra.

* Intervertebral disc space
** Clinical term: end-plates



Fig. 770 Thoracic vertebrae, Vertebrae thoracicae; AP-radiograph of the thoracic part of the vertebral column; upright position with the thorax in inspiration; the central beam is directed onto the sixth thoracic vertebra.

* Intervertebral disc space

Lumbar part of the vertebral column, radiography



Fig. 771 Lumbar vertebrae, Vertebrae lumbales;
lateral radiograph of the lumbar part of the vertebral column; upright position;
the central beam is directed onto the second lumbar vertebra. In this case, the anterior edges of the lower lumbar vertebrae are oblique due to pathological alterations.

* Intervertebral disc space
** Region of the vertebral arch between the superior and inferior articular processes ("isthmus" = interarticular portion)
*** The marks indicate the position of the twelfth rib, which is poorly visible in this copy of the radiograph.

Lumbar part of the vertebral column, radiography



Fig. 772 Lumbar vertebrae, Vertebrae lumbales;
AP-radiograph of the lumbar part of the vertebral column and the sacrum;
upright position; the central beam is directed onto the second lumbar vertebra.
* Intervertebral disc space



Fig. 773 Craniocervical junctions and the upper cervical part of the vertebral column; ventral view.



Fig. 774 Craniocervical junctions; dorsal view.

Craniocervical junctions



Fig. 775 Craniocervical junctions, deep ligaments; dorsal view.



Fig. 776 Craniocervical junctions, deep ligaments; dorsal view.



Fig. 777 Craniocervical junctions, deep ligaments; dorsal view.
The alar ligaments frequently extend to the lateral masses of the atlas.



Fig. 778 Craniocervical junctions; median section.





Fig. 779 Craniocervical junctions; superior view.



Fig. 780 Craniocervical junctions; AP-radiograph.

Ligaments of the vertebral column



Fig. 781 Ligaments of the vertebral column; exemplified by the lower thoracic part of the vertebral column; ventral view.



Fig. 782 Ligaments of the vertebral column; exemplified by the lower thoracic and the upper lumbar part of the vertebral column; dorsal view.



Fig. 783 Costovertebral joints, Articulationes costovertebrales; cross-section at the level of the lower part of the joint at a head of a rib; superior view.



Fig. 784 Ligaments of the vertebral column and the costovertebral joints, Articulationes costovertebrales; lateral parts of the anterior longitudinal ligament have been removed;
viewed from the left.



Fig. 785 Junctions between the vertebral arches; ventral view.

Costovertebral joints



Fig. 786 Costovertebral joints, Articulationes costovertebrales;
ventral view from the left.



Fig. 787 Ligaments of the vertebral arches and the costovertebral joints, Articulationes costovertebrales;
dorsal view.

* The median portion of the thoracolumbar fascia is referred to as supraspinous ligament.

Ligaments of the lumbar part of the vertebral column



Fig. 788 Ligaments of the lumbar part of the vertebral column.



Fig. 789 Vertebral joints of the lumbar part of the vertebral column, Articulationes zygapophysiales lumbales; dorsal view from the right.

Reinforcement of the interarticular joints by strong and transverse fibres (Ligg. transversa) is restricted to the lumbar part of the vertebral column.

Intervertebral discs





Fig. 790a, b Intervertebral discs, Disci intervertebrales.
a Cervical intervertebral discs, Disci intervertebrales cervicales; frontal section
b Lumbar intervertebral disc, Discus intervertebralis lumbalis; median section

* Hyaline cartilaginous covering of the end-plates of the vertebral bodies reflecting the non-ossified portion of the epiphyses

** In the first decade of life, so-called uncovertebral clefts develop in the lateral zones of the cervical intervertebral discs, which can progress further towards the middle in the following decades.



Fig. 791 Lumbar intervertebral disc, Discus intervertebralis lumbalis; superior view.



Fig. 792 Lumbar parts of the vertebral column; magnetic resonance tomographic image (MRI); median section.



Fig. 793 Cervical motion segment; schematic median section.

* Also: rim of vertebral body

** Hyaline cartilaginous covering of the end-plate of the vertebral body reflecting the non-ossified portion of the epiphysis



Fig. 794 Lumbar motion segment; schematic median section.

* Also: rim of vertebral body

** Hyaline cartilaginous covering of the end-plate of the vertebral body reflecting the non-ossified portion of the epiphysis

Superficial muscles of the back



→ T 17, T 18

Fig. 795 Muscles of the back, Mm. dorsi; superficial layer of the trunk-arm and trunk-shoulder girdle muscles.



Fig. 796 Muscles of the back, Mm. dorsi; deep layer of the trunk-arm and trunk-shoulder girdle muscles.

* The so-called thoracolumbar fascia forms a dense aponeurosis.

→ T 17–T 19

Deep muscles of the back



→ T 20a, b

Fig. 797 Muscles of the back, Mm. dorsi; superficial layer of the deep (autochthonous) muscles.





Fig. 798 Muscles of the back, Mm. dorsi;
cross-section at the level of the second lumbar vertebra;
inferior view.

The deep (autochthonous) muscles of the back lie within an osteofibrous tube formed by dorsal parts of the vertebrae and the surrounding aponeurotic thoracolumbar fascia. The muscles are divided into a lateral * and a medial ** tract.



Fig. 799 Muscles of the back, Mm. dorsi;
computed tomographic cross-section (CT) at the level of the intervertebral disc between the third and the fourth lumbar vertebra;
inferior view.

* Calcification or ossification frequently occurs at the sites of insertion of the ligamenta flava, even in younger individuals.

** Adipose tissue deposits

Deep muscles of the back



Fig. 800 Deep (autochthonous) muscles of the back; diagram of the different groups of muscles.
The deep (autochthonous) muscles of the back can be divided into a longitudinal and an oblique system, as well as into a medial and a lateral tract.

* Spinotransverse

** Transversospinal



Fig. 801 Muscles of the back, Mm. dorsi, and suboccipital muscles, Mm. suboccipitales.

→ T 20b, c

Deep muscles of the back



→ T 20 a, b

Fig. 802 Muscles of the back, Mm. dorsi; deepest layer in the region of the lower thoracic and the lumbar part of the vertebral column.



Fig. 803 Muscles of the back, Mm. dorsi, and suboccipital muscles, Mm. suboccipitales.

→ T 20 b, c



Fig. 804 Suboccipital muscles, Mm. suboccipitales.

→ T 20 c

Suboccipital muscles



→ T 20a, b

Fig. 805 Muscles of the back, Mm. dorsi, and muscles of the neck, Mm. colli.



→ T 20c

Fig. 806 Suboccipital muscles, Mm. suboccipitales.

I = Posterior tubercle of the atlas
II = Spinous process of the axis



Fig. 807 Cervical part of the vertebral column; computed tomographic cross-section (CT) at the level of the intervertebral disc between the fourth and the fifth cervical vertebra.

* Endotracheal tube and endoscope



Fig. 808 Cervical part of the vertebral column; computed tomographic cross-section (CT) at the level of the fifth and the sixth cervical vertebra.



Fig. 809 Lumbar part of the vertebral column; computed tomographic cross-section (CT) at the level of the intervertebral disc between the second and the third lumbar vertebra.



Fig. 810 Lumbar part of the vertebral column; computed tomographic cross-section (CT) at the level of the pedicles of the third lumbar vertebra.

Cutaneous innervation



Fig. 811 Segmental cutaneous innervation (dermatomes) and cutaneous nerves of the back.



813
814

Fig. 812 Vessels and nerves of the back.
* Vessels and nerves of the triangular space
** Vessels and nerves of the quadrangular space

Vessels and nerves of the posterior cervical region



814
812

Fig. 813 Vessels and nerves of the occipital region, Regio occipitalis, the posterior cervical region, Regio cervicalis posterior, and the upper back.

Vessels and nerves of the posterior cervical region



Fig. 814 Vessels and nerves of the occipital region, Regio occipitalis, and the posterior cervical region, Regio cervicalis posterior.

815
813

Cervical part of the vertebral canal



814

Fig. 815 Nerves of the posterior cervical region, Regio cervicalis posterior, and the vertebral artery, A. vertebralis.



Fig. 816 Vessels and nerves of the deep posterior cervical region, Regio cervicalis posterior, and the content of the vertebral canal.



Fig. 817 Spinal nerve, N. spinalis; in the thoracic region.



Fig. 818 Vessels and nerves of the lumbar part of the vertebral canal, Regio lumbalis.

Veins and nerves of the lumbar part of the vertebral column



Fig. 819 Veins of the vertebral canal, Canalis vertebralis; venous plexus.



Fig. 820 Nerves of the vertebral column, Columna vertebralis; somatic and autonomous innervation.



Fig. 821 Content of the vertebral canal, Canalis vertebralis; cross-section at the level of the fifth cervical vertebra.

→ 599 ff



Fig. 822 Content of the vertebral canal, Canalis vertebralis; cross-section at the level of the third lumbar vertebra.

→ 599 ff

Lumbar and sacral puncture



Fig. 823 Lumbar puncture to obtain cerebrospinal fluid, Liquor cerebrospinalis, or for lumbal anaesthesia; position of the patient.



→ 818

Fig. 824 Lumbar puncture; guidance of the puncture needle.



Fig. 825 Sacral puncture; guidance of the puncture needle.



Fig. 826 Content of the vertebral canal, Canalis vertebralis; thoracic portion after stepwise exposure; ventral view.

→ 599 ff



Fig. 827 a, b Intervertebral foramina, Foramina intervertebralia; lumbar part of the vertebral column.

a Viewed from the left
b Sagittal section

Surface anatomy



Fig. 828 Surface anatomy of the thoracic and abdominal wall of a young woman.



Fig. 829 Surface anatomy of the thoracic and abdominal wall of a young man.



Fig. 830 Projection of the skeleton onto the thoracic and abdominal wall.

Skeleton of the trunk



Fig. 831 Thoracic cage, Cavea thoracis, and left shoulder girdle, Cingulum pectorale.



Fig. 832 Ribs, Costae; I.–III. rib, superior view; VIII. rib, inferior view.

Sternum



Fig. 833 Sternum, Sternum.



Fig. 834 Sternum, Sternum.



Fig. 835 Sternum, Sternum, and costal cartilages, Cartilagines costales; section.



Fig. 836 Sternoclavicular joint, Articulatio sternoclavicularis.

Muscles of the thoracic and the abdominal wall



→ T 15, T 17, T 18, T 26

Fig. 837 Muscles of the thoracic and the abdominal wall, Mm. thoracis et abdominis.



Fig. 838 Muscles of the thoracic and the abdominal wall; superficial layer.

→ T 15, T 17, T 18, T 26

Muscles of the thoracic and the abdominal wall



→ T 13, T 15, T 26

Fig. 839 Muscles of the thoracic wall, Mm. thoracis.



→ T 13

Fig. 840 Muscles of the thoracic wall, Mm. thoracis; frontal section.
* Position of the needle for puncture of the pleural cavity (pleurocentesis)



Fig. 841 Thoracic cage, Cavea thoracis; posterior wall.

→ T 11–T 13



Fig. 842 Thoracic cage, Cavea thoracis; anterior wall.

→ T 13

Abdominal muscles



→ T 13–T 15, T 17, T 26

Fig. 843 Muscles of the abdominal wall, Mm. abdominis; superficial and middle layer.



Fig. 844 Muscles of the abdominal wall, Mm. abdominis; middle layer.

→ T 13–T 15, T 17

Abdominal muscles



→ T 13–T 15

Fig. 845 Muscles of the abdominal wall, Mm. abdominis; deep layer.



Fig. 846 Internal surface of the anterior abdominal wall.

→ T 14, T 15, T 21

Inguinal canal



Fig. 847 Superficial inguinal ring, Anulus inguinalis superficialis.



Fig. 848 Walls of the inguinal canal.



Fig. 849 Diagram of the inguinal canal.
The inguinal canal, the spermatic cord and the scrotum are illustrated in the same plane for didactical reasons.



Fig. 850 Diagram of inguinal hernias;
left side: lateral, indirect hernia;
right side: medial, direct hernia.

* Hernial sac with an intestinal loop
** Peritoneal space
*** Newly formed peritoneal hernial sac

Diaphragm and posterior abdominal wall



→ T 15, T 16, T 21, T 42

Fig. 851 Diaphragm, Diaphragma, and muscles of the abdominal wall, Mm. abdominis.



Fig. 852 Diaphragm, Diaphragma.

→ T 21



Fig. 853 Diaphragm, Diaphragma, with apertures and muscles of the posterior abdominal wall.

* Clinical term: BOCHDALEK's triangle

** Psoas arcade

→ T 21

Breast



Fig. 854 Breast, Mamma.



Fig. 855 Breast, Mamma.



Fig. 856 Breast, Mamma, of a pregnant woman.



Fig. 857 Breast, Mamma, of a pregnant woman.



Fig. 858 Radiograph of the breast, mammography, of a 47-year-old woman.



Fig. 859 Lymphatic drainage of the breast and location of the regional lymph nodes.

Vessels and nerves of the wall of the trunk



Fig. 860 Vessels and nerves of the thoracic and the abdominal wall.
* Clinical term: internal mammary artery



Fig. 861 Segmental sensory innervation of the anterior thoracic and abdominal wall (dermatomes).



Fig. 862 Segmental sensory innervation of the anterior thoracic and abdominal wall.
Regions to which pain from diseased viscera (zones of HEAD) is referred to are indicated in grey.

Lumbosacral plexus



→ 47, T 40

Fig. 863 Lumbosacral plexus, Plexus lumbosacralis.



Fig. 864 Vessels of the anterior thoracic and abdominal wall.



Fig. 865 Diagram of the collateral circulation via the internal thoracic artery, A. thoracica interna, the superior epigastric artery, A. epigastrica superior, and the inferior epigastric artery, A. epigastrica inferior, in the case of aortic occlusion; viewed from the right.

Abdominal wall of a neonate



Fig. 866 Anterior abdominal wall of a neonate.



Fig. 867 Anterior abdominal wall of a neonate.



Fig. 868 Anterior abdominal wall, internal aspect.

Abdominal wall, sections



→ T 14–T 16, T 20 a, b, T 42

Fig. 869 a–c Muscles of the abdominal wall, Mm. abdominis; horizontal sections.



Fig. 870 Muscles of the abdominal wall, Mm. abdominis; frontal section.



Fig. 871 a, b Muscles of the abdominal wall, Mm. abdominis; computed tomographic cross-sections (CT).

Heart





Abb. 872 Heart, Cor.

Pericardium fibrosum
Pericardium serosum
Lamina parietalis
} Pericardium

Lamina visceralis = Epicardium



Fig. 873 Heart, Cor, and great vessels; dorsal view.

* Arrow in the transverse pericardial sinus
** Double arrows in the oblique pericardial sinus
*** Remnants of the foetal ductus arteriosus (BOTALLO's ligament)

Myocardium



Fig. 874 Myocardium, Myocardium; ventral view.



Fig. 875 Myocardium, Myocardium; viewed from the apex of the heart.



Fig. 876 Myocardium, Myocardium; dorsoinferior view.



Fig. 877 Valves of the heart, Valvae cordis; superior view.



Fig. 878 Contour of the heart, valves of the heart, and points of auscultation projected onto the anterior thoracic wall (circles). The heart sounds and (where applicable) the heart murmurs are spreading in the direction of the arrows.

Internal spaces of the heart



Fig. 879 Right atrium, Atrium cordis dextrum, and right ventricle, Ventriculus cordis dexter; ventral view.



Fig. 880 Right atrium, Atrium cordis dextrum, of a neonate; ventral view from the right.



Fig. 881 Left atrium, Atrium cordis sinistrum, and left ventricle, Ventriculus cordis sinister; lateral view.



Fig. 882 Left ventricle, Ventriculus cordis sinister, and ascending aorta, Aorta ascendens; position of the cusps of the aortic valve (arrows) in their position at the filling phase of the heart (diastole); lateral view.

Internal spaces of the heart



Fig. 883 Left and right ventricle, Ventriculus cordis sinister et dexter.



Fig. 884 Left and right ventricle, Ventriculus cordis sinister et dexter; superior view.



Fig. 885 Left ventricle, Ventriculus cordis sinister; ventrosuperior view from the left.



Fig. 886 Right ventricle, Ventriculus cordis dexter; dorsal view.

Conducting system



Fig. 887 Right atrium, Atrium cordis dextrum, and right ventricle, Ventriculus cordis dexter; conducting system, Complexus stimulans cordis [Systema conducente cordis], highlighted in yellow; ventral view.



Fig. 888 Left ventricle, Ventriculus cordis sinister; conducting system, Complexus stimulans cordis [Systema conducente cordis], highlighted in yellow; ventral view from the left.



Fig. 889 Innervation of the heart, diagram; sympathetic trunk: green; vagus nerve: purple.
* The branches beyond the cardiac plexus contain sympathetic and vagal fibres.

→ 956



Fig. 890 Conducting system of heart, Complexus stimulans cordis [Systema conducente cordis]; section at the plane of the axis of the heart.

Coronary arteries



Fig. 891 Coronary arteries, Aa. coronariae; ventral view. Arrow in the transverse pericardial sinus



Fig. 892 Coronary arteries, Aa. coronariae; superior view.



Fig. 893 Variability of the coronary arteries. The conus arteriosus branch arises from the aorta as an independent artery (~37%).



Fig. 894 Veins of the heart, Vv. cordis.



Fig. 895 a–c Variations in the arterial supply of the posterior aspect of the heart; dorsal view.

a Balanced coronary artery supply
b Dominant left coronary artery supply
c Dominant right coronary artery supply

Coronary arteries, variability of supply



Fig. 896 Coronary arteries, Aa. coronariae; posterior arterial branches are shown in a lighter colour; the posterior interventricular branch arises from the right coronary artery (balanced coronary artery supply); ventral view.



Fig. 897 Coronary arteries, Aa. coronariae; the posterior interventricular branch arises from the left coronary artery (dominant left coronary artery supply); ventral view.



Fig. 898 Coronary arteries, Aa. coronariae; the posterior wall of the ventricles is mainly supplied by branches of the right coronary artery (dominant right coronary artery supply); ventral view.



Fig. 899 Left coronary artery, A. coronaria sinistra; coronary angiography; the beam is directed obliquely from right anterior to left posterior (RAO).



Fig. 900 Left coronary artery, A. coronaria sinistra; coronary angiography; the beam is directed obliquely from left anterior to right posterior (LAO).



Fig. 901 Right coronary artery, A. coronaria dextra; coronary angiography; the beam is directed obliquely from left anterior to right posterior (LAO).

Heart situs



Fig. 902 Position of the heart in the thorax, Situs cordis.



Fig. 903 Position of the heart; inspiration.



Fig. 904 Position of the heart; expiration.



Fig. 905 Pericardium, Pericardium.



Fig. 906 Openings of the large veins into the heart; ventral view.

Trachea and bronchi





Right main bronchus, Bronchus principalis dexter

Bronchus lobaris superior dexter
1 = Bronchus segmentalis apicalis [B I]
2 = Bronchus segmentalis posterior [B II]
3 = Bronchus segmentalis anterior [B III]

Bronchus lobaris medius dexter
4 = Bronchus segmentalis lateralis [B IV]
5 = Bronchus segmentalis medialis [B V]

Bronchus lobaris inferior dexter
6 = Bronchus segmentalis superior [B VI]
7 = Bronchus segmentalis basalis medialis [B VII]
8 = Bronchus segmentalis basalis anterior [B VIII]
9 = ronchus segmentalis basalis lateralis [B IX]
10 = ronchus segmentalis basalis posterior [B X]

Left main bronchus, Bronchus principalis sinister

Bronchus lobaris superior sinister
1,2 = Bronchus segmentalis apicoposterior [B I + II]
3 = Bronchus segmentalis anterior [B III]
4 = Bronchus lingularis superior [B IV]
5 = Bronchus lingularis inferior [B V]

Bronchus lobaris inferior sinister
6 = Bronchus segmentalis superior [B VI]
7 = Bronchus segmentalis basalis medialis [B VII]
8 = Bronchus segmentalis basalis anterior [B VIII]
9 = Bronchus segmentalis basalis lateralis [B IX]
10 = Bronchus segmentalis basalis posterior [B X]

Fig. 907 Larynx, Larynx; trachea, Trachea, and bronchi, Bronchi.



Fig. 908 Projection of the trachea and the bronchi onto the anterior thoracic wall.



Fig. 909 Larynx, Larynx; trachea, Trachea, and bronchi, Bronchi; dorsal view.



Fig. 910 Trachea, Trachea; cross-section; microscopic enlargement at low magnification.

Segmental bronchi



Fig. 911 Bronchi, Bronchi;
the lobar and the segmental bronchi have been projected onto the lungs and are illustrated with different colours; ventral view.
Numbers indicate the segmental bronchi (see p. 90).



Fig. 912 Bronchi, Bronchi;
AP-radiograph; bronchography of the left lung.
* Bronchography catheter in the trachea

Lungs



Fig. 913 Right lung, Pulmo dexter; lateral view.



Fig. 914 Left lung, Pulmo sinister; lateral view.



Fig. 915 Right lung, Pulmo dexter; sagittal section at the plane of the hilum of the lung; medial view.



Fig. 916 Left lung, Pulmo sinister; sagittal section at the plane of the hilum of the lung; medial view.

Bronchopulmonary segments



Pulmo dexter

Pulmo dexter, Lobus superior

- Segmentum apicale [S I]
- Segmentum posterius [S II]
- Segmentum anterius [S III]

Pulmo dexter, Lobus medius

- Segmentum laterale [S IV]
- Segmentum mediale [S V]

Pulmo dexter, Lobus inferior

- Segmentum superius [S VI]
- Segmentum basale mediale [cardiacum] [S VII]*
- Segmentum basale anterius [S VIII]
- Segmentum basale laterale [S IX]
- Segmentum basale posterius [S X]

* This segment is generally not an independent segment, but rather fused with the anterior basal segment [S VIII].

Fig. 917 Right lung, Pulmo dexter; bronchopulmonary segments, Segmenta bronchopulmonalia; lateral view.

Pulmo sinister

Pulmo sinister, Lobus superior

- Segmentum apicoposterius [S I + II]
- Segmentum anterius [S III]
- Segmentum lingulare superius [S IV]
- Segmentum lingulare inferius [S V]

Pulmo sinister, Lobus inferior

- Segmentum superius [S VI]
- Segmentum basale mediale [cardiacum] [S VII]*
- Segmentum basale anterius [S VIII]
- Segmentum basale laterale [S IX]
- Segmentum basale posterius [S X]

* This segment is generally not an independent segment, but rather fused with the anterior basal segment [S VIII].



Fig. 918 Left lung, Pulmo sinister; bronchopulmonary segments, Segmenta bronchopulmonalia; lateral view.



Fig. 919 Right lung, Pulmo dexter; bronchopulmonary segments, Segmenta bronchopulmonalia; medial view.



Fig. 920 Left lung, Pulmo sinister; bronchopulmonary segments, Segmenta bronchopulmonalia; medial view.

Lungs, structure



Fig. 921 Lungs, Pulmones, and heart, Cor; the arteries, the veins and the bronchi have been dissected up to the external surface of the lungs.



Fig. 922 Lung, Pulmo; overview of the lung structure; microscopic enlargement at low magnification.



Fig. 923 Vessels of the left lung, Pulmo sinister; dorsal view.



Fig. 924 Bronchi, Bronchi; arterial supply.

Vessels of the lung, radiography



Fig. 925 Arteries of the right lung,
Aa. pulmonales dextrae;
AP-radiograph (pulmonary angiography); after injection of a contrast medium into the right ventricle;
ventral view.
The numbers designate the segmental arteries (compare p. 96).



Fig. 926 Veins of the right lung,
Vv. pulmonales dextrae;
AP-radiograph (return via pulmonary veins of the contrast medium injected into the right ventricle);
ventral view.
Numbers indicate the segmental veins (compare p. 96).



Fig. 927 Thoracic cage, Cavea thoracis, and thoracic viscera; PA-radiograph of a 27-year-old male (thorax radiograph).



Fig. 928 Outline of the heart shadow in the radiograph;
Dt = Transverse diameter, ab + cd = 13–14 cm
L = Longitudinal axis of the heart (from the superior border of the right atrial contour to the apex of the heart) = 15–16 cm
M = Median plane of the body

Bronchoscopy



Fig. 929 Bronchi, Bronchi;
bronchoscopy of a healthy individual displaying the tracheal bifurcation and the carina of trachea.



Fig. 930 Bronchi, Bronchi;
bronchoscopic view of the left segmental bronchi.



Fig. 931 Projection of the pulmonary and the pleural borders onto the anterior thoracic wall.



Fig. 932 Projection of the pulmonary and the pleural borders onto the posterior thoracic wall.

Borders of the lungs – line
Borders of the pleura – dashed line



Fig. 933 Projection of the pulmonary and the pleural borders onto the lateral thoracic wall.



Fig. 934 Projection of the pulmonary and the pleural borders onto the lateral thoracic wall.

Oesophagus



Fig. 935 Oesophagus, Oesophagus; trachea, Trachea, and the thoracic aorta, Pars thoracica aortae; ventral view.



Fig. 936 Oesophagus, Oesophagus; trachea, Trachea, and the thoracic aorta, Pars thoracica aortae; viewed from the right.



Fig. 937 Oesophagus, Oesophagus, and the thoracic aorta, Pars thoracica aortae; dorsal view.



Fig. 938 Oesophagus, Oesophagus; microscopic enlargement at low magnification.

Vessels of the oesophagus



Fig. 939 Oesophagus, Oesophagus; supplying arteries; ventral view.



Fig. 940 Veins of the oesophagus, Vv. oesophageae.

→ 1077, 1078



→ 1078

Fig. 941 Veins of the oesophagus, Vv. oesophageae; magnification of Fig. 940; demonstration of anastomoses between branches of the hepatic portal vein and the superior vena cava.



Fig. 942 Thoracic lymph nodes; Nodi lymphoidei thoracis; dorsal view.

Oesophagus, radiography and oesophagoscopy



Fig. 943 a, b Oesophagus, Oesophagus; radiograph after swallowing of a contrast medium.
a Right anterior oblique position, oblique diameter I (beam directed from left anterior to right posterior)
b Left anterior oblique position, oblique diameter II (beam directed from right anterior to left posterior)

* Oesophageal constriction caused by the aortic arch
** Retrocardial portion of the oesophagus
*** Oesophageal constriction at the junction of the pharynx with the oesophagus



Fig. 944 Oesophagus, Oesophagus; oesophagoscopy (technical details see Fig. 986); superior view.



Fig. 945 Oesophagus, Oesophagus; constrictions.
* Upper oesophageal sphincter
** Narrowing caused by the aortic arch
*** Diaphragmatic constriction



Fig. 946 Oesophagus, Oesophagus; typical location and frequency of diverticula.
* Cervical diverticulum (ZENKER's diverticulum), (~70%)
** Mid-oesophageal diverticulum (~22%)
*** Epiphrenic diverticulum (~8%)

Thymus





→ 1049

Fig. 947 Thymus, Thymus; pericardium, Pericardium, and lungs, Pulmones, of an adolescent.
Note the size of the adolescent thymus. In older individuals, the thymic tissue is almost entirely replaced by adipose tissue.



Fig. 948 Thymus, Thymus, of an adolescent.

→ 1049

Pleural cavity



Fig. 949 Pleural cavity, Cavitas pleuralis, and mediastinum, Mediastinum, of an adolescent;
lateral thoracic wall and right lung have been removed; viewed from the right.
The x indicate the areas at the root of the lung and the pulmonary ligament, respectively, where the visceral pleura folds back into the parietal pleura.



Fig. 950 Pleural cavity, Cavitas pleuralis, and mediastinum, Mediastinum, of an adolescent;
viewed from the left.
In older individuals, the thymic tissue is almost entirely replaced by adipose tissue.

Aortic arch



Fig. 951 Heart, Cor, and aortic arch, Arcus aortae,

with the origins of the major arteries;
ventral view.



1 Arcus aortae
2 A. thoracica interna
3 Truncus brachiocephalicus
4 A. subclavia dextra
5 A. carotis communis dextra
6 A. vertebralis dextra
7 Rima glottidis
8 A. vertebralis sinistra
9 Trachea
10 A. subclavia sinistra
11 A. carotis communis sinistra
12 Pars descendens aortae
13 Cor
14 Valva aortae
15 Pars ascendens aortae

Fig. 952 Aortic arch, Arcus aortae, and branches;
AP-radiograph after injection of a contrast medium into the aortic bulb;
ventral view.
* Catheter





Fig. 953 a–e Variability of the origin of the major arteries from the aortic arch, Arcus aortae.
a "Textbook case"
b Common origin of the brachiocephalic trunk and the left common carotid artery
c Common trunk for the brachiocephalic trunk and the left common carotid artery
d Left vertebral artery branching off the aortic arch independently
e Right subclavian artery as the ultimate branch of the aortic arch
This abnormal artery frequently passes behind the oesophagus to the right, which can result in swallowing difficulties (Dysphagia lusoria).



Fig. 954 Thoracic and abdominal aorta, Pars thoracica aortae et Pars abdominalis aortae.

Posterior mediastinum



Fig. 955 Vessels and nerves of the posterior mediastinum, Mediastinum posterius.



Fig. 956 Autonomous nervous system of the thoracic cavity, Pars thoracica autonomica.

→ 49, 50

Phrenic nerve



Fig. 957 Thoracic viscera of an adult; the anterior thoracic wall has been removed, and the right and the left lung have been sectioned in the frontal plane; ventral view.



Fig. 958 Course of the phrenic nerve, N. phrenicus.



Fig. 959 Organisation of the mediastinum, Mediastinum.



Fig. 960 Thoracic cavity, Cavitas thoracis, and mediastinum, Mediastinum; median sagittal section through the neck and the thorax; lateral view from the right.

Thorax, frontal sections



Fig. 961 Thoracic cavity, Cavitas thoracis.



Fig. 962 Thoracic cavity, Cavitas thoracis; magnetic resonance tomographic image (MRI); frontal section at the level of the superior vena cava; ventral view.



Fig. 963 Diaphragm, Diaphragma, and oesophagus, Oesophagus, with its transition into the stomach, Gaster; frontal section through the lower part of the thoracic cavity and the upper part of the abdominal cavity.

→ 1187



Fig. 964 Thoracic cavity, Cavitas thoracis; magnetic resonance tomographic image (MRI); frontal section at the level of the aortic valve; ventral view.

Thorax, frontal sections



Fig. 965 Thoracic cavity, Cavitas thoracis; axilla, Axilla, and shoulder joint, Articulatio humeri.



Fig. 966 Thoracic cavity, Cavitas thoracis; axilla, Axilla, and shoulder joint, Articulatio humeri.



Fig. 967 a, b Thoracic cavity, Cavitas thoracis; axilla, Axilla, and shoulder joint, Articulatio humeri.

a Right part of the body
b Left part of the body

→ 294

Thorax, transverse sections



Fig. 968 Thoracic cavity, Cavitas thoracis.



Fig. 969 Thoracic cavity, Cavitas thoracis.



Fig. 970 Thoracic cavity, Cavitas thoracis.



Fig. 971 Thoracic cavity, Cavitas thoracis; computed tomographic cross-section (CT).

Thorax, transverse sections



Fig. 972 Thoracic cavity, Cavitas thoracis.



Fig. 973 Heart, Cor;
ultrasound image viewed from the oesophagus (transoesophageal echocardiography).



Fig. 974 Thoracic cavity, Cavitas thoracis.



Fig. 975 Thoracic cavity, Cavitas thoracis.

Stomach





Fig. 976 Stomach, Gaster; ventral view.



Fig. 977 Stomach, Gaster; external muscle layers; ventral view.



Fig. 978 Stomach, Gaster; the peritoneum has been removed to demonstrate the internal muscle layers; ventral view.

Stomach, structure



Fig. 979 Stomach, Gaster, and duodenum, Duodenum; ventral view.



Fig. 980 Diagram of the stomach wall; the layers of the wall have been removed stepwise; microscopic enlargement at low magnification.



Fig. 981 Blood vessels of the stomach, Gaster; ventral view.



Fig. 982 Stomach, Gaster, and liver, Hepar, with lymph nodes, Nodi lymphoidei; ventral view.

Stomach, radiography



1 = Oesophagus with contrast medium. At the transition (1a) into the fundus of stomach, the grooves between the folds appear as dark striations.
2 = Fundus of stomach with air bubble
3 = Body of stomach
3a = Lesser curvature
3b = Greater curvature. In the linings of the latter, notches corresponding to the contour of the mucous are visible.
4 = Peristaltic constriction at the angular notch
5 = Pyloric part prior to the progression of a portion of the stomach's content
6 = Ampulla of duodenum
7 = Descending part of the duodenum with circular folds
8 = Jejunum
9 = Left "dome" of diaphragm
10 = Left colic flexure (filled with air)

Fig. 983 Stomach, Gaster, and duodenum, Duodenum; AP-radiograph after oral administration of a contrast medium; upright position; ventral view.



Fig. 984 Stomach, Gaster; projection of a "normal" stomach onto the anterior abdominal wall; upright position.



Fig. 985 Stomach, Gaster; projection of a "long" stomach onto the anterior abdominal wall; upright position.



Fig. 986 Technical procedure of oesophagoscopy and gastroscopy.
* Gastroscope
** The gastroscope's tip located in the body of stomach (compare Fig. 987 a)
*** The gastroscope's tip located in the pyloric antrum (compare Fig. 987 b)



Fig. 987 a, b Stomach, Gaster; endoscopic image of the stomach (gastroscopy); superior view.

a View of the body of stomach, Corpus gastricum, with distinct longitudinal folds of the mucosa (Plica gastricae)
b View of the pyloric antrum, Antrum pyloricum, predominantly covered with smooth mucosa

Duodenum



Fig. 988 Duodenum, Duodenum.



Fig. 989 Duodenum, Duodenum, and pancreas, Pancreas; projected onto the anterior abdominal wall.



Fig. 990 Duodenum, Duodenum; ventral view.
* Clinical term: muscle of TREITZ
** Clinical term: papilla of VATER



Fig. 991 Duodenum, Duodenum; AP-radiograph after oral administration of a contrast medium; upright position; ventral view.



Fig. 992 Duodenum, Duodenum; endoscopic image.

Small intestine, structure





Fig. 993 Superior part of the duodenum, Pars superior duodeni; layers of the wall.
* Clinical term: BRUNNER's glands



Fig. 994 Small intestine, Intestinum tenue; cross-section through the upper small intestine.



Fig. 995 Small intestine, Intestinum tenue; layers of the wall; microscopic enlargement at low magnification.



Fig. 996 Jejunum, Jejunum.



Fig. 997 Ileum, Ileum.



Fig. 998 Terminal ileum, Pars terminalis ilei.
* Clinical term: PEYER's patches



Fig. 999 MECKEL's diverticulum (Diverticulum ilei). This vestige of the yolk stalk (Ductus omphaloentericus) occurs in 1–3% of cases. It is located 30–70 cm oral to the ileocaecal junction opposite to the attachment of the mesentery.

Large intestine



Fig. 1000 Large intestine, Intestinum crassum; ventral view.



Fig. 1001 Large intestine, Intestinum crassum; projected onto the anterior abdominal wall.
The positions of the transverse colon and of the sigmoid colon are highly variable (compare Fig. 1010).



Fig. 1002 Caecum, Caecum; vermiform appendix, Appendix vermiformis, and terminal ileum, Pars terminalis ilei; dorsal view.



Fig. 1003 Ascending colon, Colon ascendens; caecum, Caecum, and vermiform appendix, Appendix vermiformis; ventral view.
* Clinical term: PEYER's patches



Fig. 1004 Transverse colon, Colon transversum; ventroinferior view.



Fig. 1005 Colon, Colon; layers of the wall; microscopic enlargement at low magnification.

Vermiform appendix



Fig. 1006 Vermiform appendix, Appendix vermiformis; variations in its position; ventral view.



Fig. 1007 a–d Vermiform appendix, Appendix vermiformis; variations in its position.
a Descending into the lesser pelvis
b Behind the caecum
c In front of the ileum
d Behind the ileum



Fig. 1008 Caecum, Caecum, and vermiform appendix, Appendix vermiformis; projection onto the anterior abdominal wall.
* Clinical term: v. LANZ's point, at the right third of a line connecting the anterior superior iliac spines, locates the tip of the vermiform appendix
** Clinical term: McBURNEY's point, at the outer third of a line connecting the right anterior superior iliac spine and the umbilicus, locates the base of the vermiform appendix



Fig. 1009 Colon, Colon, and rectum, Rectum; AP-radiograph after filling with a contrast medium and air (double contrast method).



Fig. 1010 a–d Transverse colon, Colon transversum; common variations in its position; ventral view.

a Normal position
b Twisted
c U-shaped
d V-shaped



Fig. 1011 Ascending colon, Colon ascendens; endoscopic image after passage of the rectum, the sigmoid colon, the descending colon and the transverse colon (colonoscopy).

Liver



Fig. 1012 Liver, Hepar; ventral view.



Fig. 1013 Liver, Hepar, and porta hepatis, Porta hepatis; dorsal view.

* Borders of the superior recess of the omental bursa



Fig. 1014 Liver, Hepar; superior view.



Fig. 1015 Liver, Hepar, of a foetus; oxygen content of the blood is shown by different colours and the direction of flow by arrows; dorsal view.

* Also: Ductus ARANTII

Liver puncture



Fig. 1016 Liver, Hepar; projection onto the anterior abdominal wall; intermediate position between inspiration and expiration.
* Position of the needle for liver puncture



Fig. 1017 Layers of the abdominal wall and the liver, Hepar; frontal section.
* Position of the needle for liver puncture



Fig. 1018 a–d Liver, Hepar; variations in shape.
a Small left lobe of liver, Lobus hepatis sinister
b Tongue-shaped process of the right lobe of liver, Lobus hepatis dexter
c Bridged flat shape
d Diaphragmatic furrows



Fig. 1019 Liver, Hepar;
sagittal section through the right lobe of liver to demonstrate the branches of the hepatic veins and the hepatic portal vein.
* Intrahepatic branches of the hepatic veins
** Intrahepatic branches of the hepatic portal vein and the hepatic artery



Fig. 1020 Liver, Hepar;
microscopic structure (microscopic enlargement at low magnification);
trias of GLISSON (consists of a branch of the hepatic portal vein, a branch of the hepatic artery proper and a small bile duct), and a small liver lobule surrounding a central vein.

Blood vessels of the liver



Fig. 1021 Liver, Hepar, and hepatic veins, Vv. hepaticae; ventral view.



Fig. 1022 Liver, Hepar, and the hepatic portal vein, V. portae hepatis; ventral view.



Fig. 1023 Liver, Hepar;
the segments of the lobes are shown in different colours;
ventral view.



Fig. 1024 Liver, Hepar;
the segments of the lobes are shown in different colours;
dorsal view.

Liver, ultrasound



→ 1021

Fig. 1025 Hepatic veins, Vv. hepaticae;
ultrasound image showing the opening of the hepatic veins into the inferior vena cava;
inferior view.
* Abdominal wall



→ 1022

Fig. 1026 Hepatic portal vein, V. portae hepatis;
ultrasound image showing the division of the hepatic portal vein into the main branches;
inferior view.
* Abdominal wall



Fig. 1027 Liver, Hepar, and gallbladder, Vesica biliaris [fellea]; laparoscopy;
oblique inferior view from the left.



Fig. 1028 Gallbladder, Vesica biliaris [fellea], and liver, Hepar; laparoscopy;
ventral view.

Gallbladder and bile duct system



Fig. 1029 Gallbladder, Vesica biliaris [fellea], and bile duct system; ventral view.
* Clinical term: HEISTER's valve
** Clinical term: tubercle of VATER



Fig. 1030 a–c Variations in the bile duct system: common hepatic duct, Ductus hepaticus communis, and bile duct, Ductus choledochus.
a High union of the common hepatic duct, Ductus hepaticus communis, and the cystic duct, Ductus cysticus
b Low union of the common hepatic duct, Ductus hepaticus communis, and the cystic duct, Ductus cysticus
c Low union after the cystic duct, Ductus cysticus, crosses over the common hepatic duct, Ductus hepaticus communis



Fig. 1031 Bile duct system;
AP-radiograph after administration of a contrast medium;
upright position;
ventral view.



Fig. 1032 Gallbladder, Vesica biliaris [fellea], and bile duct system;
AP-radiograph after administration of a contrast medium;
upright position;
ventral view.

Pancreas



Fig. 1033 Retroperitoneal organs and vessels of the upper abdomen; ventral view.



Fig. 1034 Duodenum, Duodenum, and pancreas, Pancreas; ventral view.

* Clinical term: duct of WIRSUNG
** Clinical term: SANTORINI's duct
*** Clinical term: tubercle of VATER
**** Clinical term: muscle of TREITZ



Fig. 1035 Duodenum, Duodenum, and pancreas, Pancreas; dorsal view.



Fig. 1036 a–f Variations in the opening of the bile duct, Ductus choledochus, and the pancreatic duct, Ductus pancreaticus.
a Long common part
b Ampullary enlargement at the terminal part
c Short common part
d Separate opening
e Single opening with a septum dividing the common duct
f Accessory duct, Ductus pancreaticus accessorius



Fig. 1037 Pancreas, Pancreas; microscopic enlargement at low magnification.

Pancreas, imaging



Fig. 1038 Pancreatic duct, Ductus pancreaticus; bile duct, Ductus choledochus, and gallbladder, Vesica biliaris [fellea];
AP-radiograph; supine position; after endoscopic intubation of the common excretory duct of the liver and the pancreas, as well as injection of a contrast medium; ventral view.
Clinical term: ERCP (endoscopic retrograde cholangio-pancreaticography).



Fig. 1039 Pancreas, Pancreas;
ultrasound image showing the pancreas and adjacent large vessels in deep inspiration; oblique inferior view.
* Abdominal wall



Fig. 1040 Pancreas, Pancreas;
the gastrocolic ligament of the greater omentum has been dissected to expose the omental bursa;
photograph taken during surgery.
* Dissection line at the greater omentum



Fig. 1041 Pancreas, Pancreas;
computed tomographic horizontal section (CT); inferior view.
In this subject, the coeliac trunk is located remarkably low.
* Air bubble in the stomach

Spleen



Fig. 1042 Spleen, Splen [Lien]; ventromedial view.



Fig. 1043 Spleen, Splen [Lien]; superolateral view.



Fig. 1044 Spleen, Splen [Lien]; cross-section through the splenic hilum; superomedial view.



Fig. 1045 Development of the peritoneal cavity, Cavitas peritonealis, and the visceral relationships;
schematic median section;
lateral view.



Fig. 1046 Development of the peritoneal cavity, Cavitas peritonealis;
schematic horizontal section;
superior view.



Fig. 1047 Development of the peritoneal cavity, Cavitas peritonealis, and the visceral relationships of the female;
final state of the peritoneal cavity with adhesion of the greater omentum to the transverse colon;
schematic median section;
lateral view.
The omental bursa corresponds with the peritoneal cavity via the omental foramen.
* Clinical term: pouch of DOUGLAS

Greater omentum



→ 846

Fig. 1048 Position of the abdominal viscera, Situs viscerum, and greater omentum, Omentum majus.



Fig. 1049 Position of the abdominal viscera of the neonate.

Abdominal viscera in the upper abdomen



Fig. 1050 Position of the abdominal viscera in the upper abdomen; ventral view.
* Also: foramen of WINSLOW



Fig. 1051 Stomach, Gaster, and greater omentum, Omentum majus; photograph taken during surgery; ventral view.
* Surgical drape
** Surgical retractor



Fig. 1052 Abdominal viscera in the upper abdomen; parts of the lesser omentum have been removed to expose the omental bursa and the pancreas; oblique view from superior.



Fig. 1053 Omental bursa, Bursa omentalis; after dissection of the gastrocolic ligament; oblique view from inferior.

Abdominal viscera in the lower abdomen



Fig. 1054 Position of the abdominal viscera, Situs viscerum.



Fig. 1055 Small intestine, Intestinum tenue, and large intestine, Intestinum crassum.

Abdominal viscera in the lower abdomen



Fig. 1056 Small intestine, Intestinum tenue, and large intestine, Intestinum crassum.



Fig. 1057 Mesentery, Mesenterium, and large intestine, Intestinum crassum.

Mesenteric root and retroperitoneal space



Fig. 1058 Position of the abdominal viscera, Situs viscerum, and the omental bursa, Bursa omentalis.
An arrow indicates the omental [epiploic] foramen.



Fig. 1059 Posterior wall of the peritoneal cavity, Cavitas peritonealis, and spleen, Splen [Lien], of the female. Sites of attachment of the ascending and descending colon are indicated (✦).

* Clinical term: pouch of DOUGLAS

** Clinical term: round ligament

Abdominal arteries, overview



Fig. 1060 Arteries of the abdominal viscera; semi-schematic; possible anastomoses are indicated (○).



Fig. 1061 Hepatic portal vein, V. portae hepatis, with tributaries; semi-schematic.

Coeliac trunk



Fig. 1062 a–f Variability of the arterial blood supply of the liver.
- **a** "Textbook case"
- **b** The superior mesenteric artery, A. mesenterica superior, participates in the supply of the right lobe of the liver.
- **c** The common hepatic artery, A. hepatica communis, arises from the superior mesenteric artery, A. mesenterica superior.
- **d** The left lobe of the liver is supplied by the left gastric artery, A. gastrica sinistra.
- **e** A branch of the left gastric artery, A. gastrica sinistra, supplies the left lobe of the liver along with the left branch of the hepatic artery proper, A. hepatica propria.
- **f** An accessory branch of the hepatic artery proper, A. hepatica propria, supplies the lesser curvature of the stomach.



Fig. 1063 a–e Variability of the arterial blood supply of the stomach.
- **a** "Textbook case", closed arterial arcade supplying both the lesser and the greater curvature
- **b** The left gastric artery, A. gastrica sinistra, participates in the supply of the left lobe of the liver.
- **c** Anastomosis between the right and left gastro-omental arteries, Aa. gastroomentales dextra et sinistra, at the greater curvature
- **d** Absence of an anastomosis between the right and left gastro-omental arteries, Aa. gastroomentales dextra et sinistra, at the greater curvature
- **e** An accessory posterior gastric artery, A. gastrica posterior, arising from the splenic artery, A. splenica, supplies the posterior wall of the stomach.



Fig. 1064 Arteries of the stomach, Gaster; the spleen, Splen [Lien], and the liver, Hepar;
AP-radiograph after selective injection of a contrast medium into the coeliac trunk (coeliacography) with concomitant visualization of the renal pelvis due to the renal excretion of the medium; ventral view.

* Catheter loop in the aorta
** Catheter in the aorta



Fig. 1065 Common hepatic artery, A. hepatica communis;
AP-radiograph after selective injection of a contrast medium into the common hepatic artery;
ventral view.

* Branches to the left lobe of the liver replacing a single left branch of the hepatic artery proper
** An accessory branch of the hepatic artery to the lesser curvature of the stomach
*** Catheter in the aorta

Vessels of the upper abdomen



Fig. 1066 Vessels of the upper abdomen.



Fig. 1067 Vessels of the upper abdomen.

Superior mesenteric artery



Fig. 1068 a–d Variability of the branches of the superior mesenteric artery, A. mesenterica superior, supplying the large intestine.

a "Textbook case", the ascending and the transverse colon, Colon ascendens et transversum, are supplied by three branches.

b The ileocolic artery, A. ileocolica, and the right colic artery, A. colica dextra, form a common trunk.

c Only two branches arise from the mesenteric artery while the right colic artery, A. colica dextra, is absent.

d Duplication of the right colic artery, A. colica dextra



Fig. 1069 Superior mesenteric artery, A. mesenterica superior; AP-radiograph after injection of a contrast medium into the origin of the superior mesenteric artery; ventral view.

* Catheter in the aorta
** Catheter in the common iliac artery



Fig. 1070 Superior mesenteric artery and vein, A. et V. mesenterica superior. The arteries supplying the small intestine form a series of arcades.

Inferior mesenteric artery



Fig. 1071 a–c Variations in the branches of the inferior mesenteric artery, A. mesenterica inferior.
a The inferior mesenteric artery, A. mesenterica inferior, divides into three parts supplying the descending and sigmoid colon, as well as the rectum.
b An accessory middle colic artery, A. colica media, arises from the inferior mesenteric artery, A. mesenterica inferior.
c An accessory middle colic artery, A. colica media, arises from the left colic artery, A. colica sinistra.
* Clinical term: SUDECK's point



Fig. 1072 Inferior mesenteric artery, A. mesenterica inferior; AP-radiograph after selective injection of a contrast medium into the origin of the inferior mesenteric artery; ventral view.
* Catheter in the aorta



Fig. 1073 Inferior mesenteric artery and vein, A. et V. mesenterica inferior.

Coeliac trunk





Fig. 1074 Abdominal aorta, Pars abdominalis aortae, with the coeliac trunk, Truncus coeliacus, and the superior mesenteric artery, A. mesenterica superior; ultrasound image; almost in the sagittal plane.
* Abdominal wall



Fig. 1075 a–e Variability of the coeliac trunk, Truncus coeliacus.
a "Textbook case", division of the trunk into three branches
b Division of the trunk into four branches
c Development of a hepatosplenic trunk, Truncus hepatosplenicus
d Development of a gastrosplenic trunk, Truncus gastrosplenicus
e Development of a gastrosplenic trunk, Truncus gastrosplenicus, and a hepatomesenteric trunk, Truncus hepatomesentericus



Fig. 1076 Vessels of the retroperitoneal space, Situs retroperitonealis, of the male.

Hepatic portal vein, overview



Fig. 1077 Hepatic portal vein, V. portae hepatis; ventral view.



Fig. 1078 Hepatic portal vein, V. portae hepatis, and inferior vena cava, V. cava inferior; semi-schematic; tributaries of the inferior vena cava in blue; tributaries of the hepatic portal vein in purple; potential portocaval anastomoses are indicated by circles.





Fig. 1079 Kidney, Ren [Nephros], and suprarenal gland, Glandula suprarenalis; ventral view (left).



Fig. 1080 Kidney, Ren [Nephros], and suprarenal gland, Glandula suprarenalis; ventral view (right).



Fig. 1081 Kidney, Ren [Nephros]; oblique and vertical hemisection displaying the renal cortex, the medulla and the pelvis; ventral view (left). Arrows point from the pyramids to the calyces.



Fig. 1082 Projection of the kidneys onto the back.



Fig. 1083 Kidney, Ren [Nephros]; oblique and vertical hemisection; ventral view (left).



Fig. 1084 Kidney, Ren [Nephros]; transverse section through the renal sinus; inferior view (left).



Fig. 1085 Kidney, Ren [Nephros]; microscopic section of the cortex (100×).



Fig. 1086 Kidney, Ren [Nephros], and the suprarenal gland, Glandula suprarenalis, of a ~ 5-month-old foetus.



Fig. 1087 Kidney, Ren [Nephros]; dorsal view.
The lower poles of both kidneys are fused (= horseshoe kidney).

Renal pelvis and calyces



Fig. 1088 a, b Renal pelvis, Pelvis renalis; corrosion casts; ventral view (left).
a Dendritic type
b Ampullar type



Fig. 1089 Kidney, Ren [Nephros]; renal pelvis, Pelvis renalis, and ureter, Ureter; AP-radiograph after retrograde injection of a contrast medium via both ureters.



Fig. 1090 a, b Renal segments, Segmenta renalia.

a Ventral view (right)
b Dorsal view (right)



Fig. 1091 Kidney, Ren; areas of contact with adjacent organs; ventral view.

Areas of contact of the kidneys

- Glandulae suprarenales
- Hepar
- Duodenum, Pars descendens
- Colon, Flexura dextra
- Jejunum
- Gaster
- Splen [Lien]
- Pancreas
- Colon descendens



Fig. 1092 Kidney, Ren [Nephros];
ultrasound image;
the transducer is directed from ventroinferior to dorsosuperior;
lateral view (right).
* Abdominal wall



→ 1196

Fig. 1093 Kidney, Ren [Nephros];
puncture of the right kidney;
computed tomographic cross-section (CT);
inferior view.
* Guidance of a needle for kidney biopsy



Fig. 1094 Suprarenal gland, Glandula suprarenalis; ventral view (right).



Fig. 1095 a–d Variability of the suprarenal arteries, Aa. suprarenales, and course of the suprarenal veins, Vv. suprarenales.
a Arterial supply by three arteries ("textbook case")
b Arterial supply without a branch from the renal artery, A. renalis
c Arterial supply without a direct branch from the aorta, Aorta
d Course of the suprarenal veins, Vv. suprarenales



Fig. 1096 Suprarenal gland, Glandula suprarenalis; sagittal section; lateral view (right).



Fig. 1097 Suprarenal gland, Glandula suprarenalis; microscopic enlargement at low magnification.



Fig. 1098 Urinary bladder, Vesica urinaria; prostate, Prostata, and urethra, Urethra; ventral view.



Fig. 1099 Urinary bladder, Vesica urinaria, almost completely voided.



Fig. 1100 Urinary bladder, Vesica urinaria, filled. In this situation, the bladder can be punctured from just above the pubic bone without passing through the peritoneal cavity.
* Puncture needle



Fig. 1101 Urinary bladder, Vesica urinaria; ductus deferentes, Ductus deferentes; seminal vesicles, Glandulae vesiculosae, and prostate, Prostata; dorsal view.





Fig. 1102 a, b Urinary bladder, Vesica urinaria; endoscopic view of the opening of the ureter (cystoscopy).
a Open ostium of the ureter with a peristaltic wave transporting urine into the bladder
b Closed ostium of the ureter

Fig. 1103 Urinary bladder, Vesica urinaria; endoscopic view of the mucosa in the body of the bladder (cystoscopy); inferior view.



→ 1119

Fig. 1104 Male urinary and genital organs;
Organa urogenitalia masculina;
diagram of the development: the parts that degenerate are displayed in pale pink, and the position of the testis prior to its descent is indicated by the dashed line;
viewed from the right.

Epididymis = genital part of the mesonephros;
Paradidymis = remnants of tubules of the mesonephros

* WOLFFian duct
** MÜLLERian duct
*** Junction of the MÜLLERian ducts, Ductus paramesonephrici



Fig. 1105 Testis, Testis [Orchis], and epididymis, Epididymis; viewed from the right.



Fig. 1106 Testis, Testis [Orchis], and epididymis, Epididymis; viewed from the right.



Fig. 1107 Blood vessels of the testis, Testis, the epididymis, Epididymis, and the spermatic cord, Funiculus spermaticus; viewed from the right.



Fig. 1108 Testis, Testis; epididymis, Epididymis, and ductus deferens (vas deferens), Ductus deferens; lateral view.



Fig. 1109 Testis, Testis; epididymis, Epididymis, and scrotum, Scrotum; superior view.
* Also: Epiorchium
** Also: Periorchium



Fig. 1110 Ductus deferentes, Ductus deferentes, seminal vesicles, Glandulae vesiculosae, and prostate, Prostata; superior view.



Fig. 1111 Ductus deferentes, Ductus deferentes, and seminal vesicles, Glandulae vesiculosae; AP-radiograph after injection of a contrast medium via the ejaculatory ducts; ventral view.

Though this technique demonstrates size and position of the seminal vesicles, it is no more applied under clinical circumstances.



Fig. 1112 Urinary bladder, Vesica urinaria; prostate, Prostata; ductus deferens (vas deferens), Ductus deferens, and seminal vesicle, Glandula vesiculosa; viewed from the right.



Fig. 1113 Urinary bladder, Vesica urinaria; ductus deferentes, Ductus deferentes; seminal vesicles, Glandulae vesiculosae; prostate, Prostata, and male urethra, Urethra masculina; dorsal view.



Fig. 1114 Male external genitalia, Organa genitalia masculina externa.



Fig. 1115 Spermatic cord, Funiculus spermaticus; frontal section; ventral view (left, 250%).



→ 1199

Fig. 1116 Urinary bladder, Vesica urinaria; prostate, Prostata, and male urethra, Urethra masculina; ventral view.



Fig. 1117 a–d Penis, Penis; cross-sections; planes indicated in Fig. 1118; ventral view.

a Cross-section through the middle of the shaft
b Cross-section at the level of the proximal part of the glans penis
c Cross-section through the middle of the glans penis
d Cross-section at the level of the distal part of the glans penis



Fig. 1118 Penis, Penis, with glans penis, Glans penis, and prepuce, Preputium penis.

* Level of section of Fig. 1117a
** Level of section of Fig. 1117b
*** Level of section of Fig. 1117c
**** Level of section of Fig. 1117d

Female urinary and genital organs, overview



→ 1104

Fig. 1119 Female urinary and genital organs, Organa urogenitalia feminina; diagram of the development: the parts that degenerate are displayed in pale pink, and the position of the ovary prior to its descent is indicated by the dashed line; ventral view.

Epoophoron = genital part of the mesonephros;
Paroophoron = remnants of the tubules of the mesonephros

* WOLFFian duct
** MÜLLERian duct
*** BARTHOLIN's glands



Fig. 1120 Uterus, Uterus; ovary, Ovar, and uterine tube, Tuba uterina; dorsal view.

* Clinical term: cardinal ligament
** Clinical term: sacrouterine ligament



Fig. 1121 Uterus, Uterus; ovary, Ovar, and uterine tube, Tuba uterina; dorsal view.

* Clinical term: GRAAFian follicle
** Stalked hydatid
*** Clinical term: round ligament



Fig. 1122 Ovary, Ovar;
microscopic enlargement at low magnification.



Fig. 1123 Uterus, Uterus, and vagina, Vagina;
viewed from the right.



Fig. 1124 Mucosa of the uterus, Uterus;
early phase of proliferation.



Fig. 1125 Mucosa of the uterus, Uterus;
late phase of proliferation with excretory glands.



Fig. 1126 Uterus, Uterus, and vagina, Vagina; normal angles between the vagina, the cervix, and the body of uterus;
viewed from the right.
* Longitudinal axis of the vagina
** Longitudinal axis of the cervix of uterus
*** Longitudinal axis of the body of uterus
Angle between vagina and cervix of uterus = version
Angle between cervix and body of uterus = flexion
Normal topographical situation of the uterus: anteversion, anteflexion
Relation to the median plane = position



Fig. 1127 a–c Uterus, Uterus, and vagina, Vagina.
a Anteversion, anteflexion = normal position
b Anteversion, but no anteflexion
c Retroversion, retroflexion



Fig. 1128 Uterus, Uterus, and uterine tube, Tuba uterina; AP-radiograph after injection of to contrast medium into the cervix of uterus (hysterosalpingography);
uterus in dextroposition;
ventral view.
This formerly clinically applied technique allows to determine the position of the organs and the patency of the tubes.
K = tube adaptor for injection of the contrast medium



Fig. 1129 Arteries of the female internal genitalia, Organa genitalia feminina interna; dorsal view.
* Clinical term: round ligament



Fig. 1130 a–f Varibility of the arterial supply of the female internal genitalia; dorsal view.
a Supply of the uterus, Uterus ("textbook case")
b, c, d Supply of the ovaries, Ovaria (b, "textbook case")
e, f Supply of the fundus of uterus, Fundus uteri (e, "textbook case")



Fig. 1131 Uterus, Uterus;
uterine ligaments and connective tissue spaces; semi-schematic transverse section at the level of the cervix of uterus;
superior view.

* Clinical term: Paraproctium
** Clinical term: Parametrium
(existence of this structure is controversial)
*** Clinical term: Paracystium



Fig. 1132 a, b Vaginal portion of the cervix of uterus, Portio vaginalis cervicis.
a Photograph of the cervix of a young nulliparous woman
b Photograph of the cervix of a young woman who has given birth to two children

For the inspection of the vaginal portion of the cervix the normally slit-like vagina is spreaded by means of a bivalve speculum (*);
inferior view.



Fig. 1133 Uterus, Uterus, with an embryo; ultrasound image taken in the eighth week of pregnancy; lateral view (right).

The embryo is immersed in the amniotic fluid of the chorionic cavity.

Fig. 1134 Uterus, Uterus, with a foetus; ultrasound image taken in the 28th week of pregnancy; lateral view (left).

Ultrasound examination allows the visualization of movements of the extremities and the opening of the mouth.



Fig. 1135 Hand, Manus, of a foetus; ultrasound image taken in the 18th week of pregnancy; lateral view.

Details such as fingers can be observed.



Fig. 1136 Uterus, Uterus, with a foetus; the pelvis has been sectioned in the median plane.
* Mucous plug (of KRISTELLER) in the cervical canal of the uterus
** Clinical term: pouch of DOUGLAS
*** Clinical term: vesicovaginal septum
**** Clinical term: cave of RETZIUS



Fig. 1137 Uterus, Uterus; position of the fundus of uterus during pregnancy. Numbers refer to the end of the respective month of pregnancy (= 28 days).

a



b



Fig. 1138 a, b Placenta, Placenta, and umbilical cord, Funiculus umbilicalis.
a View of the foetal surface
b View of the maternal surface of a parturient placenta



Fig. 1139 Female internal genitalia, Organa genitalia feminina interna; ventral view.
* Clinical term: infundibulopelvic ligament



Fig. 1140 Female internal genitalia, Organa genitalia feminina interna; surgical exposure in a young woman; ovaries displaced both medially and superiorly by compresses (*) in the pouch of DOUGLAS; ventrosuperior view.
** Swab



Fig. 1141 Abdominal ostium of the uterine tube, Ostium abdominale tubae uterinae; surgical exposure in a young woman; the pelvic cavity is filled with saline to demonstrate the fimbria; dorsosuperior view.
* Plastic tray to support the uterine tube



Fig. 1142 Rectum, Rectum; viewed from the left.



Fig. 1143 Rectum, Rectum; blood supply; the mucosa and the pararectal adipose tissues have been partly removed; openings into the hepatic portal vein in purple.



Fig. 1144 Rectum, Rectum, and anus, Anus; ventral view.
* Haemorrhoidal node
** KOHLRAUSCH's fold



Fig. 1145 Rectum, Rectum;
endoscopic image of the anal canal with six enlarged nodes of the cavernous rectal body, haemorrhoids; superior view.
* Colonoscope
** Three haemorrhoidal nodes





Fig. 1146 Rectum, Rectum;
endoscopic image of the rectal ampulla (rectoscopy); inferior view.



Fig. 1147 Rectal arteries, Aa. rectales; dorsal view.

* Clinical term: SUDECK's point (from this point on there are no further anastomoses with the sigmoid arteries)



Fig. 1148 Rectal veins, Vv. rectales; diagram with parts of the pelvis and the pelvic diaphragm; dorsal view.
There are numerous connections between the veins draining into the hepatic portal vein (superior rectal vein) and those draining into the inferior vena cava (middle and inferior rectal veins). They form the portocaval anastomoses, which are of particular clinical importance.

→ 1077, 1078



Fig. 1149 Rectum, Rectum; schematic overview of the innervation; ventral view.
Green = sympathetic nervous system
Purple = parasympathetic nervous system



Fig. 1150 Rectum, Rectum;
lateral radiograph in voluntary closure of the anus after filling with a contrast medium (defaecography).

The transitional zone between the anus and the rectum (arrow) is located at the level of the tip of the coccyx (triangle). The angle between the axes of the anus and the rectum (∡) is approximately 90° and depends upon the curvature of the levator ani muscle (puborectal muscle). Scale in cm.



Fig. 1151 Rectum, Rectum;
lateral radiograph of the defaecation after filling with a contrast medium (defaecography).

In comparison to Fig. 1150, the anorectal transitional zone has descended and the angle (∡) increased to 137° due to the relaxation of the curvature of the levator ani muscle. As the bending acts like a valve, the elongation results in an unimpeded pressure of the faeces on the anal canal leading to defaecation.

Arterial supply of the kidney



Fig. 1152 Kidney, Ren;
AP-radiograph after intravenous injection of a contrast medium which is excreted via the kidneys to demonstrate the renal pelvis and the ureters (intravenous pyelography); concomitant visualization of the arteries by injection of a contrast medium into the renal artery through a catheter* introduced into the aorta (arteriography).



Fig. 1153 a–d Variations in the arterial supply of the kidney.
a One renal artery, A. renalis, with a branch to the superior pole
b Two renal arteries to the renal hilum
c Two renal arteries, one of which supplies the superior pole
d Two renal arteries, one of which supplies the inferior pole





Fig. 1154 Position of the retroperitoneal structures, Situs retroperitonealis, of the male.
Whereas the left testicular vein opens into the left renal vein, the right testicular vein drains directly into the inferior vena cava. The same applies to the ovarian veins.

Fig. 1155 a, b Variability of the course of the testicular arteries, Aa. testiculares.
a "Textbook case"
b Both testicular arteries, Aa. testiculares, branch off cranially to the renal veins, Vv. renales; the right artery passes posterior to the inferior vena cava, V. cava inferior, and the left artery passes anterior to the left renal vein, V. renalis sinistra.

Lymphatics of the retroperitoneal space



→ 1389

Fig. 1156 Lymph nodes, Nodi lymphoidei, and lymphatics, Vasa lymphatica, of the posterior abdominal wall and the inguinal region;
ventral view.

The numbers X and XII indicate the respective ribs.

* Clinical term: "horizontal chain", draining the lower abdominal wall, the gluteal region, the perineum and the external genital



Fig. 1157 Lymphatics, Vasa lymphatica, and lymph nodes, Nodi lymphoidei, of the inguinal, the pelvic and the lumbar region;
AP-radiograph after bilateral injection of a contrast medium into lymphatics of the foot (lymphography).
This formerly applied technique visualizes the position and size of lymphatics and lymph nodes.



→ 47, 49, 50

Fig. 1158 Nerves of the posterior abdominal wall, the lumbosacral plexus, Plexus lumbosacralis, and the abdominal part of the autonomous nervous system, Pars abdominalis autonomica; ventral view.



Fig. 1159 Vessels and nerves of the posterior abdominal wall of the male.

* In ~10% the left renal vein passes posterior to the aorta.



Fig. 1160 Male genitalia,
Organa genitalia masculina;
diagram demonstrating the autonomous innervation of the left side;
ventral and lateral view, respectively.
Green = sympathetic nervous system
Purple = parasympathetic nervous system

* Also: erigent nerves

Surgical removal of paraaortic lymph nodes or operations at the abdominal aorta and the large arteries of the pelvis can cause damage to the sympathetic nerves with the consequence of ejaculatory impotency (Impotentia generandi). Prostate operations, in turn, may damage the parasympathetic fibres innervating the penis, thus leading to erectile impotency (Impotentia coeundi).

Innervation of the male genitalia

	Origin	Course	Organ	Function
Parasympathetic	Sacral part of the spinal cord (S2 – S4)	Ganglia pelvica, Radix parasympathica [Nn. splanchnici pelvici]	Penis Corpus cavernosum	Vasodilatation Erection
Sympathetic	Thoracic part of the spinal cord (T10 – T12)	Plexus mesenterici superior et inferior ↓ Truncus sympathicus ↓ Plexus testicularis ↓	Testis	Regulation of blood flow
	Lumbar part of the spinal cord (L1 – L2)	Plexus hypogastricus superior ↓ N. hypogastricus ↓		
		Plexus hypogastricus inferior	Glandula bulbo-urethralis	Expulsion of its fluid
			Ductus (vas) deferens	Contraction, transport of the sperms into the urethra
			Glandula vesiculosa Prostata	Expulsion of its content into the urethra
Somatomotor, somatosensory	Sacral part of the spinal cord (S2 – S4)	N. pudendus	(M. sphincter vesicae)	Closure of the urinary bladder to prevent retrograde ejaculation
			M. ischiocavernosus M. bulbospongiosus	Expulsion of the ejaculate out of the urethra
		Nn. scrotales posteriores N. dorsalis penis	Scrotal skin Penile skin	



Fig. 1161 Female genitalia, Organa genitalia feminina; diagram demonstrating the autonomous innervation; ventral view.
Green = sympathetic nervous system
Purple = parasympathetic nervous system

* Clinical term: FRANKENHÄUSER's ganglion
** Also: erigent nerves

The autonomic plexus are interspersed with ganglia. The hypogastric and uterovaginal plexus contain both sympathetic and parasympathetic fibres.

→ 49, 50

Innervation of the female genitalia

	Origin	Course	Organ	Function
Parasympathetic	Sacral part of the spinal cord (S2 – S4)	Ganglia pelvica, Radix parasympathica [Nn. splanchnici pelvici] ↓ Nn. cavernosi clitoridis	Tuba uterina Uterus Vagina Clitoris	Vasodilatation Vasodilatation Transsudation Erection
Sympathetic	Thoracic part of the spinal cord (T10 – T12)	Plexus mesentericus superior ↘ Plexus ovaricus ↗ Plexus renalis Truncus sympathicus ↓	Ovar	Vasoconstriction
	Lumbar part of the spinal cord (L1 – L2)	Plexus hypogastricus superior ↓ N. hypogastricus Plexus hypogastricus inferior ↓	-	
		Plexus uterovaginalis (FRANKENHÄUSER's ganglion)	Tuba uterina Uterus Vagina	Contraction
Somatomotor, somatosensory	Sacral part of the spinal cord (S2 – S4)	N. pudendus ↗ N. dorsalis clitoridis N. pudendus ↘ Nn. labiales posteriores	Clitoris Labia majora M. ischiocavernosus M. bulbospongiosus	 Contraction



Fig. 1162 a–d Variability of the branching pattern of the internal iliac artery, A. iliaca interna; viewed from the left.
a All branches arise from the same stem.
b The internal iliac artery, A. iliaca interna, divides into two major branches ("textbook case").
c The internal iliac artery, A. iliaca interna, divides into three major branches.
d The internal iliac artery, A. iliaca interna, divides into more than three major branches.



Fig. 1163 Internal iliac artery, A. iliaca interna, and sacral plexus, Plexus sacralis; viewed from the left.

Organs of the male pelvis



Fig. 1164 Organs of the male pelvis.
* Clinical term: COWPER's gland



Fig. 1165 Blood supply of the male pelvis.

Vessels of the female pelvis



Fig. 1166 Blood supply of the female pelvis.



Fig. 1167 Lymphatics, Vasa lymphatica, and lymph nodes, Nodi lymphoidei, of the pelvic wall of the female.

The lymph nodes are frequently much smaller than illustrated but they are always present. Tumour cells from the uterus can reach the superficial inguinal lymph nodes via lymphatics of the round ligament of uterus.

* Clinical term: pouch of DOUGLAS

** Clinical term: round ligament



Fig. 1168 a–c Variability of the origin of the obturator artery, A. obturatoria; medial view.

a Origin from the anterior branch of the internal iliac artery, A. iliaca interna ("textbook case")

b Origin as an independent branch from the internal iliac artery, A. iliaca interna

c Origin from the external iliac artery, A. iliaca externa

Only in 75% of the cases, the obturator artery originates from the trunk of the internal iliac artery.



Fig. 1169 Perineal muscles, Mm. perinei, and the pelvic diaphragm, Diaphragma pelvis, of the female; inferior view.

* Probe in the pudendal canal (ALCOCK's canal)



Fig. 1170 Pelvic diaphragm, Diaphragma pelvis, of the female; superior view.

* Clinical term: levator hiatus (Hiatus urogenitalis et ani)



Fig. 1171 Muscles of the pelvic diaphragm, Diaphragma pelvis, of the female.



→ T 22

Fig. 1172 Perineum, Perineum, and pelvic diaphragm, Diaphragma pelvis, of the male; inferior view.

* Clinical term: COWPER's gland



Fig. 1173 Perineum, Perineum; pelvic diaphragm, Diaphragma pelvis, and female external genitalia, Organa genitalia feminina externa; inferior view.
* Clinical term: BARTHOLIN's gland

→ T 22

The vaginal and the anal orifices are located in close proximity. During delivery the skin and muscles of the perineum can rupture up to the sphincter muscles of the anus (I. – III. degree perineal lacerations). These can be prevented by surgical incisions directed either laterally or in the median plane (perineal incision = lateral or medial episiotomy).



→ T 22

Fig. 1174 Urogenital diaphragm, Diaphragma urogenitale, of the male; inferior view.
* Clinical term: COWPER's gland



→ T 22

Fig. 1175 Urogenital diaphragm, Diaphragma urogenitale, of the female; inferior view.



Fig. 1176 Female external genitalia, Organa genitalia feminina externa; ventroinferior view.
* Clinical term: BARTHOLIN's gland



Fig. 1177 Female external genitalia, Organa genitalia feminina externa; inferior view.

Fig. 1178 Female external genitalia, Organa genitalia feminina externa; inferior view.

Male pelvic organs and retroperitoneal space





Fig. 1179 Pelvic diaphragm, Diaphragma pelvis; pelvic organs and anterior abdominal wall, Organa abdominis et pelvis, of the male; frontal section through the head of the femur and the urinary bladder on the left; dorsal view.



Fig. 1180 Pelvic diaphragm, Diaphragma pelvis, and pelvic organs, Organa pelvis, of the female; frontal section (on the right) combined with a median section; ventral view.



Fig. 1181 Male external genitalia,
Organa genitalia masculina externa.



Fig. 1182 Male genitalia, Organa genitalia masculina.

→ 1105, 1117



Fig. 1183 Vessels and nerves of the perineum, Regio perinealis, and the male external genitalia, Organa genitalia masculina externa.



Fig. 1184 Vessels and nerves of the perineum, Regio perinealis, and the female external genitalia, Organa genitalia feminina externa.



Fig. 1185 Abdomen, Abdomen, and pelvis, Pelvis, of the male; median section.



Fig. 1186 Abdomen, Abdomen, and pelvis, Pelvis, of the male;
sagittal section to the right of the median plane.



→ 870

Fig. 1187 Abdomen, Abdomen;
frontal section through the most anterior part of the abdominal cavity.





Fig. 1188 Abdomen, Abdomen;
frontal section demonstrating the diaphragm,
the organs of the upper abdomen and the kidneys;
dorsal view.





Fig. 1189 Abdomen, Abdomen; sagittal section through the upper abdomen at the level of the right kidney; viewed from the right.





Fig. 1190 Abdomen, Abdomen;
sagittal section through the upper abdomen at the level of the spleen;
viewed from the left.



Fig. 1191 Abdomen, Abdomen; transverse section through the upper abdomen at the level of the eleventh thoracic vertebra.



Fig. 1192 Abdomen, Abdomen; computed tomographic cross-section (CT) at the level of the tenth thoracic vertebra; inferior view.





Fig. 1193 Abdomen, Abdomen;
transverse section through the upper abdomen at the level of the intervertebral disc between the eleventh thoracic and first lumbar vertebra;
with the exception of the omental bursa highlighted in yellow green the peritoneum is coloured in blue.



Fig. 1194 Abdomen, Abdomen;
computed tomographic section (CT) at the level of the first lumbar vertebra.
The intestine is partially filled with a contrast medium.





Fig. 1195 Abdomen, Abdomen; transverse section through the upper abdomen at the level of the first lumbar vertebra.



Fig. 1196 Abdomen, Abdomen; computed tomographic section (CT) at the level of the first lumbar vertebra; inferior view.
* Intestinal gas





Fig. 1197 Abdomen, Abdomen; transverse section through the lower abdomen at the level of the third lumbar vertebra.



Fig. 1198 Abdomen, Abdomen; computed tomographic section (CT) at the level of the third lumbar vertebra; inferior view.

* Jejunum filled with a contrast medium



Fig. 1199 Pelvis, Pelvis, of the male; median section; viewed from the left.

* Clinical term: KOHLRAUSCH's fold

** Clinical term: cave of RETZIUS



Fig. 1200 Pelvis, Pelvis, of the male; magnetic resonance tomographic image (MRI); paramedian section; viewed from the left.



Fig. 1201 Pelvis, Pelvis, of the female; median section;
the intestine has been almost completely removed with some remnants of the sigmoid and rectum;
viewed from the left.

* Clinical term: vesicovaginal septum
** Clinical term: sacrouterine ligament
*** Clinical term: pouch of DOUGLAS



Fig. 1202 Pelvis, Pelvis, of the female; magnetic resonance tomographic image (MRI);
paramedian section;
viewed from the left.



Fig. 1203 Pelvis, Pelvis;
transverse section at the level of the fifth lumbar vertebra. This male specimen is different from Fig. 1191, 1193, 1195 and 1197.

As the sigmoid reaches far into the upper abdomen the figure also displays the dome of colonic flexure. The thickness of the adipose tissue layer on the gluteus medius muscle must be taken into consideration when injecting intramuscularly.



Fig. 1204 Pelvis, Pelvis;
computed tomographic cross-section (CT) at the level of the first sacral vertebra after administration of a contrast medium into the colon; supine position; inferior view.
* Calcification in the wall of the iliac artery.



Fig. 1205 Abdomen, Abdomen; volume-guided reconstruction in the frontal plane based on horizontal computed tomographic sections (CT). The section planes of organs such as the heart, the stomach, the liver and parts of the small intestine, appear granulated.

→ 1070, 1187



Fig. 1206 Pelvis, Pelvis, of the male; transverse section through the lesser pelvis.



Fig. 1207 Pelvis, Pelvis, of the male; computed tomographic cross-section (CT) through the lesser pelvis with the subject in supine position.

* Calcification in the medial part of the femoral artery



Fig. 1208 Pelvis, Pelvis, of the female; transverse section through the lesser pelvis at the level of the symphysis.





Fig. 1209 Pelvis, Pelvis, of the female; computed tomographic cross-section (CT) through the lesser pelvis with the subject in supine position.
* Remnants of contrasting intestinal contents



Fig. 1210 Pelvis, Pelvis, of the male; oblique frontal section through the lesser pelvis.
* Clinical term: ALCOCK's canal



Fig. 1211 Pelvis, Pelvis, of the male; magnetic resonance tomographic image (MRI); frontal section at the level of the hip joints; ventral view.



Fig. 1212 Pelvis, Pelvis, of the female; magnetic resonance tomographic image (MRI); frontal section at the level of the hip joints; ventral view.

When the urinary bladder is empty, the uterus lies on the apex of the bladder due to its anteflexion.

Male pelvic diaphragm, angled-frontal section



Fig. 1213 Pelvis, Pelvis, of the male; angled-frontal section through the urinary bladder.

* Clinical term: paracystium

** Clinical term: prostatic venous plexus

Female pelvic diaphragm, angled-frontal section





Fig. 1214 Pelvis, Pelvis, of the female; angled-frontal section through the urinary bladder.
* Paracystium with venous plexus

Surface anatomy



Fig. 1215 Lower limb, Membrum inferius.



Fig. 1216 Lower limb, Membrum inferius.



Fig. 1217 Lower limb, Membrum inferius; skeleton and joints.

Pelvis



Fig. 1218 Sacrum, Os sacrum, and pelvic girdle, Cingulum pelvicum.
The region cranial to the linea terminalis is referred to as the greater pelvis, Pelvis major, and the region caudal to it is known as lesser pelvis, Pelvis minor.



Fig. 1219 Pelvis, Pelvis; shape and dimensions of the pelvic inlet of the female.



Fig. 1220 Pelvis, Pelvis; shape of the pelvic inlet of the male.

k–k = Pelvic axis
a–b = Anatomical conjugate
a–e = Diagonal conjugate 12.5–13 cm
a–c = True conjugate 10.4–11 cm

h–d = Sagittal diameter of the pelvic brim 12–12.5 cm
e–g = Sagittal diameter of the pelvic constriction 11–11.5 cm
e–f = Sagittal diameter of the pelvic outlet (= Distantia pubococcygea) 9–10 cm



Fig. 1221 Pelvis, Pelvis; dimensions in the female; median section.



a–a = Crestal distance 28–29 cm*
b–b = Anterior spinal distance 25–26 cm*
c–c = Posterior spinal distance (width of the sacrum) 10 cm
* In this perspective, the crestal distance appears shorter than the anterior spinal distance.

d–d = Transverse diameter of the pelvic brim (= interacetabular line) 12–12.5 cm
e–e = Transverse diameter of the pelvic constriction (= interspinal line) 10.5 cm
f–f = Transverse diameter of the pelvic outlet (= tuberal diameter) 11–12 cm)

Fig. 1222 Pelvis, Pelvis, dimensions in the female.

Hip bone



Fig. 1223 Hip bone, Os coxae; medial view.
The thin region in the centre of the ala of ilium is characteristic for its frame-like structure.



Fig. 1224 Hip bone, Os coxae; ventral view.



Fig. 1225 Hip bone, Os coxae; dorsolateral view.



Fig. 1226 Hip bone, Os coxae; development.
* Approximately sixth year of life



Fig. 1227 Hip bone, Os coxae; stage of development in a 6-year-old child.
In the acetabulum, the three parts of the hip bone are fused to a Y-shaped cartilaginous junction. This junction ossifies at the age of 13–18 years.

Joints of the pelvic girdle



Fig. 1228 Joints of the pelvic girdle, Juncturae cinguli pelvici, and the lumbosacral joint, Articulatio lumbosacralis, of the male.





Fig. 1229 Joints of the pelvic girdle, Juncturae cinguli pelvici, and the lumbosacral joint, Articulatio lumbosacralis, of the female.



Fig. 1230 Joints of the pelvic girdle, Juncturae cinguli pelvici, and the lumbosacral joint, Articulatio lumbosacralis, of the male.





Fig. 1231 Joints of the pelvic girdle, Juncturae cinguli pelvici, and the lumbosacral joint, Articulatio lumbosacralis, of the female.

Joints of the pelvic girdle



Fig. 1232 Joints of the pelvic girdle, Juncturae cinguli pelvici, and the lumbosacral joint, Articulatio lumbosacralis, of the female.



Fig. 1233 Joints of the pelvic girdle, Juncturae cinguli pelvici, of the female.



Fig. 1234 Joints of the pelvic girdle, Juncturae cinguli pelvici, of the female.



Fig. 1235 Joints of the pelvic girdle, Juncturae cinguli pelvici, and the lumbosacral joint, Articulatio lumbosacralis, of the female; median section.

Normally, the anterior border of the lowest intervertebral disc forms the furthest projecting point of the posterior circumference of the pelvic inlet. Radiographically, the most anterior part of the sacrum is referred to as promontory.

Joints of the pelvic girdle





Fig. 1236 Sacroiliac joint, Articulatio sacroiliaca; frontal section.



Fig. 1237 Pubic symphysis, Symphysis pubica; oblique section in the direction of the longitudinal axis of the symphysis.
The interpubic disc consists of fibrous cartilage, with the exception of the articular symphysial surfaces of both pubic bones that are covered with hyaline cartilage. A longitudinal flat cleft occurs after the first decade of life (*).



Fig. 1238 Pelvis, Pelvis, of the male; AP-radiograph with the central beam directed onto the third sacral segment; upright position.

Pelvis, development



Fig. 1239 Pelvis, Pelvis, and femur, Femur; AP-radiograph of a female, premature baby (eighth month of pregnancy).

* Osseous roof of the acetabulum
** Y-shaped junction in the acetabular fossa
*** The ossification centre in the head of the femur does not appear before the third to fifth month of life.
\+ At this age, both greater trochanters only become apparent as small protuberances of the bone of the diaphysis.



Fig. 1240 Pelvis, Pelvis, and femur, Femur; AP-radiograph of a 12-month-old boy.

* Osseous roof of the acetabulum
** Y-shaped junction in the acetabular fossa
*** Ossification centre in the epiphysis of the head of the femur
\+ At this age, both greater trochanters only become apparent as small protuberances of the bone of the diaphysis.



Fig. 1241 Femur, Femur; ventral view.



Fig. 1242 Femur, Femur; dorsal view.

Femur



Fig. 1243 Femur, Femur;
proximal extremity;
dorsal view.



Fig. 1244 Femur, Femur;
variability of the angle of the neck of the femur;
dorsal view.
The angle between the neck and the shaft of the femur is referred to as neck-shaft angle. It is 150° in infants and about 126° in the adult.



Fig. 1245 Femur, Femur;
trabecular structure of the femoral bone with a large neck-shaft angle (coxa valga); section in the plane of the angle of anterior torsion (60%).
The lateral plates of the trabecular bone* ("tensile system") are only poorly developed, whereas the medial plates** ("compressive system") are much stronger.



Fig. 1246 Femur, Femur;
trabecular structure of the femoral bone with a small neck-shaft angle (coxa vara); section in the plane of the angle of anterior torsion (60%).
The lateral plates of the trabecular bone * ("tensile system") are well developed, while the medial plates ** ("compressive system") are developed less strong. As a result of the high demand of flexion, the cortex on the inner side of the femoral neck is particularly thick.



Fig. 1247 Femur, Femur;
medial view.



Fig. 1248 Femur, Femur;
variability of the angle of anterior torsion; the proximal end of the femur has been projected over the distal end; proximal view.
The angle of anterior torsion is about 30° in infants and about 14° in adults.



Fig. 1249 Femur, Femur;
cross-section through the middle of the shaft; distal view.

Hip joint



Fig. 1250 Hip joint, Articulatio coxae; ventrodistal view.



Fig. 1251 Hip joint, Articulatio coxae; dorsal view.



Fig. 1252 Hip joint, Articulatio coxae; the articular capsule has been opened and the head of the femur has been partly exarticulated; laterodistal view.



Fig. 1253 Hip joint, Articulatio coxae; aspect of the acetabulum after removal of the articular capsule and exarticulation of the head of the femur; laterodistal view.



Fig. 1254 Hip joint, Articulatio coxae; vertical section at the plane of the angle of antetorsion.
* Radiations into the iliotibial tract

Hip joint, radiography



Fig. 1255 Hip joint, Articulatio coxae; AP-radiograph; upright position on both legs.

* Clinical term: acetabular roof = tangential projection of the lunate surface
** Clinical term: edge of the acetabular roof = the most lateral projection of the acetabulum
*** Clinical term: KÖHLER's teardrop = projection of the acetabular fossa



Fig. 1256 Hip joint, Articulatio coxae; AP-radiograph; supine position with the thigh abducted and flexed (so-called LAUENSTEIN projection).

* Due to minimal absorption of X-rays by cartilage, the articular cleft appears abnormally broad.



Fig. 1257 Femur, Femur; distal extremity; lateral view.



Fig. 1258 Femur, Femur; distal extremity; distal view.



Fig. 1259 Femur, Femur; frontal section through the distal part; ventral view.

Tibia



Fig. 1260 Tibia, Tibia; ventral view.



Fig. 1261 Tibia, Tibia; lateral view.



Fig. 1262 Tibia, Tibia; dorsal view.



Fig. 1263 Tibia, Tibia, and fibula, Fibula; proximal view.

* The articular surfaces of the condyles together are referred to as superior articular surface.



Fig. 1264 Fibula, Fibula; medial view.



Fig. 1265 Fibula, Fibula; lateral view.



Fig. 1266 Tibia, Tibia, and fibula, Fibula; dorsal view.



Fig. 1267 Tibia, Tibia, and fibula, Fibula; cross-section with the interosseous membrane of leg; distal view.
* Opening for the anterior tibial artery

Patella and knee joint



Fig. 1268 Patella, Patella; ventral view.



Fig. 1269 Patella, Patella; dorsal view.



Fig. 1270 Knee joint, Articulatio genus; cross-section.
* Medial border facet



Fig. 1271 Knee joint, Articulatio genus; intact articular capsule; ventral view.



Fig. 1272 Knee joint, Articulatio genus; the anterior part of the articular capsule has been reflected downwards after dissection of the quadriceps muscle; the suprapatellar bursa has been opened; ventral view.

Knee joint



Fig. 1273 Knee joint, Articulatio genus; in 90° flexion; the articular capsule and the lateral ligaments have been removed; ventral view.

* Clinical term: ACL (= anterior cruciate ligament)
** Clinical term: PCL (= posterior cruciate ligament)



Fig. 1274 Knee joint, Articulatio genus; intact articular capsule with muscular origins; dorsal view.



Fig. 1275 Knee joint, Articulatio genus;
exposure of the cruciate ligaments and the menisci;
dorsal view.



Fig. 1276 Tibial collateral ligament, Lig. collaterale tibiale;
knee in extended position;
medial view.
Only the posterior fibres of the tibial collateral ligament are attached to the medial meniscus.

Fig. 1277 Tibial collateral ligament, Lig. collaterale tibiale;
knee in flexed position (90°);
medial view.
In the course of flexion, the posterior and proximal fibres of the tibial collateral ligament become twisted, thereby stabilizing the medial meniscus.

Knee joint



Fig. 1278 Knee joint, Articulatio genus;
synovial membrane tightly filled after injection of plastic material;
lateral view.
Subpopliteal recess not shown (compare Fig. 1279).



Fig. 1279 Knee joint, Articulatio genus;
synovial membrane thightly filled after injection of plastic material;
dorsal view.
* Clinical term: MCL (= medial collateral ligament)
** Clinical term: LCL (= lateral collateral ligament)



Fig. 1280 Menisci of the knee joint, Articulatio genus, and cruciate ligaments, Ligg. cruciata; proximal view.



Fig. 1281 Menisci of the knee joint, Articulatio genus; arterial supply; proximal view.



Fig. 1282 a, b Displacement of the menisci during flexion; lateral view.

a Extended position
b Flexed position



Fig. 1283 Displacement of the menisci during flexion; proximal view.
In flexion, both menisci are displaced posteriorly over the edges of the tibial condyles. The greater mobility of the lateral meniscus accounts for its lesser risk of damage.

Knee joint, sections



Fig. 1284 Knee joint, Articulatio genus; frontal section.



Fig. 1285 Knee joint, Articulatio genus; sagittal section through the lateral part of the joint.



Fig. 1286 Knee joint, Articulatio genus; AP-radiograph with the central beam directed onto the middle of the joint; subject reclined.



Fig. 1287 Knee joint, Articulatio genus; lateral radiograph with the central beam directed onto the middle of the joint; subject reclined.

Knee joint, arthroscopy



1 Arthroscope
2 Inlet or outlet for the rinsing fluid
3 Cold light source
4 Ocular or connector for the video system
5 Anterolateral access
6 Anteromedial access
7 Additional instrument

Fig. 1288 Accesses for arthroscopy.



Fig. 1289 a–c Knee joint, Articulatio genus; arthroscopy.
a Distal view into the femoropatellar joint
* Patellar roof ridge: ridge between the medial and lateral articular surfaces
** Clinical term: suprapatellar recess



b Medial view of the free inner edge of the lateral meniscus;
retractor (*) slightly depressing the anterior part of the meniscus



c Anterolateral view of the distal part of the anterior cruciate ligament, Lig. cruciatum anterius.
The ligament is covered by a richly vascularized synovial membrane. It is drawn medially by a retractor (*).



Fig. 1290 Knee joint, Articulatio genus; magnetic resonance tomographic image (MRI); frontal section; knee in the extended position. In this imaging technique, compact bone appears black.



Fig. 1291 Knee joint, Articulatio genus; magnetic resonance tomographic image (MRI); sagittal section through the lateral part of the joint; knee in the extended position.



Fig. 1292 a, b Knee joint, Articulatio genus; magnetic resonance tomographic image (MRI); sagittal sections to demonstrate the cruciate ligaments; knee in the extended position.

a Anterior cruciate ligament, Lig. cruciatum anterius, (ACL)
b Posterior cruciate ligament, Lig. cruciatum posterius, (PCL)
* The inhomogeneity is due to oblique sectioning of the fibre bundles.

Joints of the bones of the leg



Fig. 1293 Joints of the bones of the leg; ventral view.



Fig. 1294 Tibia, Tibia, and fibula, Fibula; distal view.

Fig. 1295 Skeleton of the foot, Ossa pedis; proximal view.





Fig. 1296 Skeleton of the foot, Ossa pedis; plantar view.

Skeleton of the foot



Fig. 1297 Skeleton of the foot, Ossa pedis; medial view.



Fig. 1298 Skeleton of the foot, Ossa pedis; lateral view.

* Also: CHOPART's joint
** Also: LISFRANC's joint



Fig. 1299 Talus, Talus; proximal view.



Fig. 1300 Talus, Talus; plantar view.



Fig. 1301 Calcaneus, Calcaneus; medial view.



Fig. 1302 Calcaneus, Calcaneus; lateral view.

Tarsal bones



Fig. 1303 Tarsal bones, Ossa tarsi, and metatarsal bones, Ossa metatarsi; proximal view.
* The cuboid bone is shown in a medial view.



Fig. 1304 a, b Skeleton of the foot, Ossa pedis; structural organisation.
a Proximal view
b Plantar view

The heads of all metatarsal bones lie in the plantar plane. The cuneiform bones, the navicular bone and the talus rest upon the lateral parts of the skeleton, especially in the posterior part of the foot. The talus is therefore situated above the calcaneus and the longitudinal arch is formed.
The wedge-shaped cross-section of the cuneiform bones and the bases of the metatarsal bones contribute to the formation of the transverse vault.



Fig. 1305 Supporting structures of the medial arch of the longitudinal vault of the foot; medial view.

The ligaments shown in this illustration are mostly directed in the longitudinal axis of the foot and support passively the longitudinal vault of the foot. The short muscles of the foot primarily support this construction.

Joints of the foot



Fig. 1306 Joints of the foot, Articulationes pedis; ligaments and tendons; medial view.



Fig. 1307 Joints of the foot, Articulationes pedis; ligaments and tendons; lateral view.
* Also: Achilles tendon



Fig. 1308 Joints of the foot, Articulationes pedis; ligaments and tendons of the posterior part of the foot.
* Also: Achilles tendon



Fig. 1309 Ankle joint, Articulatio talocruralis; proximal articular surfaces; distal view.



Fig. 1310 Joints of the foot, Articulationes pedis; plantar view.
* With the long plantar ligament the groove for the tendon of the fibularis longus muscle is enclosed in a canal.



Fig. 1311 Joints of the foot, Articulationes pedis; ligaments and tendons of the tarsus and metatarsus; plantar view.



Fig. 1312 Talotarsal joint, Articulatio talotarsalis; distal articular surfaces; proximal view.



Fig. 1313 Talotarsal joint, Articulatio talotarsalis; proximal articular surfaces; distal view.



Fig. 1314 Talotarsal joint, Articulatio talotarsalis; the talus and the lateral ligaments have been removed; lateral view.
The two arrows point out the helical distorsion of the talocalcaneal interosseous ligament.

* Tight connective tissue layer between the plantar calcaneonavicular ligament and the tibionavicular part of the deltoid ligament limiting a medially directed gliding of the head of the talus; slackening of this layer results in flattening of the longitudinal vault (flat-foot, pes valgus)

Ankle and talotarsal joint, sections



Fig. 1315 Ankle and talotarsal joint, Articulationes talocruralis et talotarsalis; frontal section.



Fig. 1316 Ankle and talotarsal joint, Articulationes talocruralis et talotarsalis; sagittal section.
* Fat pad of the heel



Fig. 1317 Ankle joint, Articulatio talocruralis; AP-radiograph with the subject reclined and the central beam directed tangentially onto the trochlea of talus.

* Clinically, the posterior border of the fibular notch is also referred to as the third malleolus.



Fig. 1318 Ankle and talotarsal joint, Articulationes talocruralis et talotarsalis; AP-radiograph with the subject reclined and the central beam directed tangentially onto the apex of the trochlea of talus.

* Due to their helical distorsion, the joint clefts are not displayed orthogonally.
** Overlapping epiphyseal lines of the tibia and fibula

→ 1428, 1429

Fasciae of the lower limb



Fig. 1319 Fascia lata, Fascia lata, and deep fascia of leg, Fascia cruris.
The transition zone between the aponeurosis of the external oblique abdominal muscle and the fascia lata is known as inguinal ligament. This ligament extends laterally from the anterior superior iliac spine to its attachment on the pubic tubercle medially.
* Openings in the fascia for the perforator veins (DODD's veins)



Fig. 1320 Fascia lata, Fascia lata, and deep fascia of leg, Fascia cruris.

Muscles of the lower limb, overview



→ T 42, T 44, T 45, T 47, T 48

Fig. 1321 Muscles of the lower limb, Mm. extremitatis inferioris; overview.



Fig. 1322 Muscles of the lower limb, Mm. extremitatis inferioris; overview.

→ T 43, T 46, T 49

Origins and insertions of the muscles of the hip, the thigh and the leg



→ T 42–T 48

Fig. 1323 Origins and insertions of the muscles at the lower lumbar vertebrae, the pelvic bones, the femur and the proximal extremities of the bones of the leg; ventral view.

Origins and insertions of the muscles of the hip, the thigh and the leg



Fig. 1324 Origins and insertions of the muscles at the pelvic bones, the femur and the proximal extremities of the bones of the leg; dorsal view.

→ T 43–T 46, T 49, T 50

Muscles of the thigh



Fig. 1325 Tensor fasciae latae muscle, M. tensor fasciae latae, and sartorius muscle, M. sartorius.

→ T 44



Fig. 1326 Quadriceps femoris muscle, M. quadriceps femoris.

→ T 44



Fig. 1327 Adductor muscles, Mm. adductores.

→ T 45



Fig. 1328 Iliopsoas muscle, M. iliopsoas.

→ T 42



Fig. 1329 Muscles of the thigh and the hip; after removal of the fascia lata except for the iliotibial tract.

* Common insertion of the sartorius, gracilis and semitendinosus muscles just below the medial condyle of the tibia (formerly known as the superficial pes anserinus, Pes anserinus superficialis)

→ T 42, T 44, T 45

Muscles of the thigh



→ T 42, T 44, T 45

Fig. 1330 Muscles of the thigh and the hip; after removal of the fascia lata, the tensor fasciae latae muscle and the sartorius muscle.

* The fourth head of the quadriceps muscle, the vastus intermedius muscle, is covered by the rectus femoris muscle.



Fig. 1331 Muscles of the thigh and the hip; deep layer after removal of the sartorius, the rectus femoris and the adductor longus muscles as well as parts of the iliopsoas muscle in the joint region.

* Common insertion of the sartorius, gracilis and semitendinosus muscles just below the medial condyle of the tibia
** The origin of the rectus femoris muscle has been flapped laterally upwards.
*** A part of the adductor longus muscle has been flapped upwards.

→ T 42, T 44, T 45

Muscles of the thigh



→ T 42, T 44, T 45

Fig. 1332 Muscles of the thigh and the hip; after almost complete removal of superficial and several deeper muscles; ventral view.



Fig. 1333 Muscles of the thigh and the hip; medial view.

→ T 22a, T 42–T 46

Muscles of the thigh and the hip



Fig. 1334 Dorsal muscles of the thigh.

→ T 46



Fig. 1335 Gluteus maximus muscle, M. gluteus maximus.

→ T 43



Fig. 1336 Gluteus medius muscle, M. gluteus medius.

→ T 43



Fig. 1337 Deep dorsal muscles of the hip.

→ T 43



Fig. 1338 Muscles of the thigh and the hip.

→ T 43



Fig. 1339 Muscles of the thigh and the hip; the gluteus maximus muscle has been sectioned; dorsal view.

* The part of the obturator internus muscle between the curved edge of the lesser sciatic notch and the insertion in the trochanteric fossa frequently consists of tendinous bands.

→ T 43, T 45, T 46

Muscles of the thigh and the hip



→ T 43, T 45, T 46

Fig. 1340 Muscles of the thigh and the hip; the gluteus maximus and medius muscles have been partially removed; dorsal view.



Fig. 1341 Muscles of the thigh and the hip; deep layer after almost complete removal of the superficial gluteal and the ischiocrural muscles; dorsal view.

→ T 43, T 45, T 46

Muscles of the thigh and the hip



→ T 43, T 44, T 46

Fig. 1342 Muscles of the thigh and the hip; after removal of the fascia lata except for the iliotibial tract.



Fig. 1343 Muscles in the region of the knee joint; medial view.

→ T 44–T 46, T 49

* Common insertion just below the medial condyle of the tibia (formerly known as superficial pes anserinus, Pes anserinus superficialis)



Fig. 1344 Muscles in the region of the knee joint; after almost complete removal of the ischiocrural muscles; dorsal view.

→ T 44–T 46, T 49

Origins and insertions of the muscles of the thigh and the ventral part of the leg



→ T 44–T 47

Fig. 1345 Origins and insertions of the muscles at the bones of the leg; ventral view.

Origins and insertions of the muscles of the thigh and the dorsal part of the leg



Fig. 1346 Origins and insertions of the muscles at the bones of the leg; dorsal view.

→ T 46, T 48–T 50

Muscles of the leg

Fig. 1347 Extensor muscles of the leg.

→ T 47





Fig. 1348 Lateral (fibular) muscles.

→ T 48



Fig. 1349 Triceps surae muscle, M. triceps surae.

→ T 49



Fig. 1350 Deep flexor muscles of the leg.

→ T 50



Fig. 1351 Muscles of the leg and the foot.

→ T 47, T 48

Muscles of the leg



→ T 49

Fig. 1352 Muscles of the leg; superficial layer.
* Also: Achilles tendon



Fig. 1353 Muscles of the leg; superficial layer after partial removal of the gastrocnemius muscle.
* Also: Achilles tendon

→ T 49

Muscles of the leg



→ T 50

Fig. 1354 Muscles of the leg; deep layer.
* Also: Achilles tendon



Fig. 1355 Muscles of the leg; deepest layer.

→ T 50

Muscles of the leg



→ T 47–T 49, T 51

Fig. 1356 Muscles of the leg and the foot.

Origins and insertions of the muscles of the the foot and the leg



Fig. 1357 Origins and insertions of the muscles at the bones of the foot; dorsal view.

→ T 47, T 48, T 51, T 53, T 54

Tendon sheaths of the foot



Fig. 1358 Tendon sheaths, Vaginae tendinum, of the foot.



Fig. 1359 Tendon sheaths, Vaginae tendinum, of the foot.



Fig. 1360 Tendon sheaths, Vaginae tendinum, of the foot.

Muscles of the foot



→ T 47, T 51, T 53

Fig. 1361 Muscles of the foot; after removal of the tendon sheaths.



Fig. 1362 Muscles of the foot; after section of the inferior extensor retinaculum.

→ T 47, T 51, T 53

Muscles of the foot



Fig. 1363 Muscles of the foot;
plantar aponeurosis, Aponeurosis plantaris.

Origins and insertions of the muscles of the foot and the leg



Fig. 1364 Origins and insertions of the muscles at the bones of the foot; plantar view.

→ T 47, T 48, T 50, T 52–T 54

Muscles of the foot



→ T 52–T 54

Fig. 1365 Muscles of the foot; the plantar aponeurosis has been removed.



Fig. 1366 Muscles of the foot; middle layer after almost complete removal of the flexor digitorum brevis muscle.

→ T 50, T 52–T 54

Muscles of the foot



→ T 50, T 52 – T 54

Fig. 1367 Muscles of the foot; deep layer.

* The site where the tendon of the flexor digitorum longus muscle crosses the tendon of the flexor hallucis longus muscle is commonly referred to as plantar chiasm, Chiasma plantare.



Fig. 1368 Dorsal interossei muscles, Mm. interossei dorsales pedis; dorsal view.

→ T 53



Fig. 1369 Plantar interossei muscles, Mm. interossei plantares; plantar view.

→ T 53

Lumbosacral plexus



→ 863

Fig. 1370 Lumbosacral and coccygeal plexus, Plexus lumbosacralis (Plexus lumbalis et sacralis) et coccygeus; segmental organisation of nerves.



Fig. 1371 Nerves of the lower limb; overview.



Fig. 1372 Nerves of the lower limb; overview.

Cutaneous innervation



Fig. 1373 Segmental cutaneous innervation (dermatomes) of the lower limb.



Fig. 1374 Segmental cutaneous innervation (dermatomes) of the lower limb.





Fig. 1375 Cutaneous nerves of the lower limb.

Fig. 1376 Cutaneous nerves of the lower limb.

Femoral and obturator nerve



Fig. 1377 Femoral and obturator nerve, Nn. femoralis et obturatorius; overview;
cutaneous innervation illustrated with purple.



Fig. 1378 Gluteal nerves, Nn. glutei, and sciatic nerve, N. ischiadicus; overview;
cutaneous innervation illustrated with purple.

Tibial and fibular nerve



Fig. 1379 Tibial nerve, N. tibialis; overview; cutaneous innervation illustrated with purple.



Fig. 1380 Fibular nerve, N. fibularis; overview; cutaneous innervation illustrated with purple.



Fig. 1381 Arteries of the lower limb; overview; ventral view.
The segment of the femoral artery between the point where the deep artery of thigh branches off and the point of entry into the adductor canal (*) is known clinically as superficial femoral artery.



Fig. 1382 Arteries of the lower limb; overview; dorsal view.

Veins and lymphatics of the lower limb



Fig. 1383 a, b Veins and lymphatics of the lower limb; overview.
a Medial view
b Dorsal view



Fig. 1384 a, b Perforating veins, Vv. perforantes; overview (according to HACH, 1986).
a Medial view
b Dorsal view



Fig. 1385 Veins of the lower limb; principle of organisation.
Disturbances of the venous flow of the lower limb, especially varicosis, are a common cause of vascular disorders.
If one of the venous systems is entirely occluded, the perforating veins may be of crucial importance to maintain the venous drainage.



Fig. 1386 Epifascial vessels and nerves of the inguinal region, Regio inguinalis, the anterior region of thigh, Regio femoris anterior, and the anterior region of knee, Regio genus anterior.

860
1391
1403

Lymphatics of the inguinal region



Fig. 1387 Superficial lymphatics, lymph nodes and major veins of the inguinal region, Regio inguinalis.



Fig. 1388 Draining areas of the lymph nodes in the inguinal region, Regio inguinalis, of the female; overview.
The arrows indicate possible directions of the flow of lymph.
* In some rare cases, the medial parts of the uterine tube and the fundus of the uterus may be drained by superficial lymph nodes in the inguinal region via the round ligament of uterus.



Fig. 1389 Saphenous opening, Hiatus saphenus, and vascular space, Lacuna vasorum; after removal of the anterior abdominal wall and the contents of the abdomen, as well as dissection of the iliac fascia and the femoral septum [CLOQUET].
* Also: ROSENMÜLLER's node
** Also: GIMBERNAT's ligament



Fig. 1390 Muscular and vascular space, Lacunae musculorum et vasorum; oblique section at the level of the inguinal ligament.

Vessels and nerves of the thigh



Fig. 1391 Vessels and nerves of the anterior region of thigh, Regio femoris anterior;
after removal of the fascia lata except for the iliotibial tract.

* Clinically the femoral artery is commonly referred to as superficial femoral artery to distinguish it from the deep artery of thigh.



Fig. 1392 Vessels and nerves of the anterior region of thigh, Regio femoris anterior;
after partial removal of the sartorius muscle and dissection of the pectineus muscle.

* The entrance into the adductor canal is formed by the vastus medialis and adductor longus muscles as well as by the anteromedial intermuscular septum spanning between them.

Vessels and nerves of the thigh



1154
1156
1159
1392 1408

Fig. 1393 Vessels and nerves of the anterior region of thigh, Regio femoris anterior; deep layer after partial removal of the sartorius and rectus femoris muscles, the pectineus and adductor longus muscles have been dissected; the adductor canal is almost entirely opened.



Fig. 1394 Arteries of the hip and the thigh; overview.

The branching pattern of these arteries varies considerably. This type of origin and arborisation of the deep artery of thigh occurs in approximately 58% of cases.



Fig. 1395 Arteries of the pelvis and the thigh; digital subtraction angiography (DSA).

Veins and nerves of the thigh



812 1398 1404

Fig. 1396 Epifascial veins and nerves of the posterior region of thigh, Regio femoris posterior, the gluteal region, Regio glutealis, and the popliteal fossa, Fossa poplitea.



Fig. 1397 Skeleton contour and sciatic nerve, N. ischiadicus, projected onto the surface of the gluteal region, Regio glutealis.



Fig. 1398 Vessels and nerves of the gluteal region, Regio glutealis, the posterior region of thigh, Regio femoris posterior, and the popliteal fossa, Fossa poplitea.

1399
1396
1411

Vessels and nerves of the thigh



1400
1398

Fig. 1399 Vessels and nerves of the gluteal region, Regio glutealis, the posterior region of thigh, Regio femoris posterior, and the popliteal fossa, Fossa poplitea; the long head of the biceps femoris muscle has been retracted laterally.
In this specimen, the medial and lateral sural cutaneous nerves branch off quite far proximally.



Fig. 1400 Vessels and nerves of the gluteal region, Regio glutealis, the posterior region of thigh, Regio femoris posterior, and the popliteal fossa, Fossa poplitea; after dissection of the gluteus maximus muscle and the long head of the biceps femoris muscle.

1399 1411 1413 1414

Vessels and nerves of the gluteal region



Fig. 1401 Vessels and nerves of the gluteal region, Regio glutealis; after dissection and partial removal of the gluteus maximus and medius muscles; the sciatic nerve has been removed after its passage through the infrapiriform foramen.



Fig. 1402 Ventrogluteal injection (according to v. HOCHSTETTER). In order to avoid the superior gluteal nerve and, in particular, the superior gluteal artery the injection is made into the triangular field formed by the two spread fingers and the iliac crest. The index finger – or when using the right hand, the middle finger – is placed on the anterior superior iliac spine and the palm of the hand on the greater trochanter. Since the injected material should be deposited within the belly of the gluteus medius muscle as far away as possible from any vessel, the needle should not cross the lines of the triangle. However, some risk remains for the nerve branch that runs from the superior gluteal nerve to the tensor fasciae latae muscle.



Fig. 1403 Epifascial veins and nerves of the leg and the foot, Regiones cruris et pedis.



Fig. 1404 Epifascial veins and nerves of the leg and the foot, Regiones cruris et pedis; the deep fascia of leg has been dissected proximally.
* Clinical term: MAY's vein

Vessels and nerves of the leg



Fig. 1405 Vessels and nerves of the popliteal fossa, Fossa poplitea;
after dissection of the deep fascia of leg and partial removal of the small saphenous vein.



1 Compartimentum cruris anterius:
A.; V. tibialis anterior
N. fibularis profundus
M. tibialis anterior
M. extensor digitorum longus
M. extensor hallucis longus
M. fibularis [peroneus] tertius

2 Compartimentum cruris laterale:
N. fibularis superficialis
M. fibularis [peroneus] longus
M. fibularis [peroneus] brevis

3 Compartimentum cruris posterius, Pars profunda:
A.; V. tibialis posterior
A.; V. fibularis
N. tibialis
M. flexor digitorum longus
M. tibialis posterior
M. flexor hallucis longus

4 Compartimentum cruris posterius, Pars superficialis:
M. triceps surae
M. plantaris

Fig. 1406 Osteofibrous tubes of the leg;
cross-section superior to the middle of the leg.
These osteofibrous tubes together with their contents are clinically known as compartments.
* Deep part of the deep fascia of leg

→ 1425



Abb. 1407 Vessels of the anterior region of leg, Regio cruris anterior;
the extensor muscles spread by retractors.

The superficial lymphatics regularly follow the major epifascial veins and converge along the great saphenous vein to the medial side of the leg. The deep lymphatics are found within the connective tissue sheaths of the deep arteries and veins of the leg.

Arteries and nerves of the leg



1391
1392
1416
1407

Fig. 1408 Arteries and nerves of the anterior region of leg, Regio cruris anterior, and the dorsum of foot, Dorsum pedis; after removal of the deep fascia of leg and dissection of the extensor digitorum longus and fibularis [peroneus] longus muscles.



Fig. 1409 Vessels and nerves of the popliteal fossa, Fossa poplitea; after removal of the fascia lata and the deep fascia of leg.



Fig. 1410 Arteries of the popliteal fossa, Fossa poplitea; demonstration of the arterial supply after partial removal of the covering muscles.
This branching pattern is found in about 90% of the cases.

Vessels and nerves of the popliteal fossa and the leg



Fig. 1411 Vessels and nerves of the popliteal fossa, Fossa poplitea, and the posterior region of leg, Regio cruris posterior; after removal of the deep fascia of leg and dissection of the gastrocnemius muscle.

Fig. 1412 a–d Variations in the branching pattern of the popliteal artery, A. poplitea.
- **a** Common trunk of the anterior and posterior tibial artery, Aa. tibiales anterior et posterior, along with the fibular artery, A. fibularis
- **b** The popliteal artery, A. poplitea, branches off proximally to the superior border of the popliteal muscle, M. popliteus.
- **c** Posterior tibial and fibular artery, Aa. tibialis posterior et fibularis, originate from a common proximal trunk.
- **d** The anterior tibial artery, A. tibialis anterior, passes ventrally to the popliteal muscle, M. popliteus.



Vessels and nerves of the popliteal fossa and the leg



Fig. 1413 Vessels and nerves of the popliteal fossa, Fossa poplitea, and the posterior region of leg, Regio cruris posterior; deep layer.



Arteries and nerves of the leg



1398
1400
1413
1421

Fig. 1414 Arteries and nerves of the popliteal fossa, Fossa poplitea, and the posterior region of leg, Regio cruris posterior; deepest layer.



1403 1416

Fig. 1415 Epifascial veins and nerves of the dorsum of foot, Dorsum pedis.

Arteries and nerves of the foot



1408
1415

Fig. 1416 Arteries and nerves of the dorsum of foot.
Dorsum pedis.



Fig. 1417 Arteries of the sole of foot, Planta; overview.



Fig. 1418 a–d Variations in the arterial supply of the sole of foot, Planta.

a Dorsalis pedis artery, A. dorsalis pedis, as major supply for the deep plantar arch, Arcus plantaris profundus

b Posterior tibial artery, A. tibialis posterior, as major supply for the deep plantar arch, Arcus plantaris profundus

c The fifth toe and lateral parts of the fourth toe are supplied by the posterior tibial artery, A. tibialis posterior, whereas the remaining medially located toes receive their arterial supply through the dorsalis pedis artery, A. dorsalis pedis.

d The fifth and fourth toe as well as lateral parts of the third toe are supplied by the posterior tibial artery, A. tibialis posterior, whereas all remaining medially located toes receive their arterial supply through the dorsalis pedis artery, A. dorsalis pedis.

Arteries and nerves of the foot



1411 1420

Fig. 1419 Arteries and nerves of the sole of foot, Planta.





Fig. 1420 Arteries and nerves of the sole of foot, Planta; deep layer.

* The distal extension of the medial retromalleolar space beneath the abductor hallucis muscle is also known as tarsal tunnel.

Arteries of the foot



1414
1420

Fig. 1421 Arteries and nerves of the sole of foot, Planta; deepest layer.
* Anastomosis with the dorsalis pedis artery



Fig. 1422 Compartments of the foot; stepwise section.
* Space of the interossei muscles
** Lateral compartment
*** Medial compartment
**** Intermediate compartment

Thigh, oblique section



Fig. 1423 Thigh, Femur; oblique section through the hip joint.



Fig. 1424 Thigh, Femur;
cross-section through the middle of the thigh.



Fig. 1425 Leg, Crus;
cross-section through the middle of the leg.

→ 1406

Leg and foot, sections



Fig. 1426 Leg, Crus;
cross-section just above the ankle joint.



Fig. 1427 Foot, Pes;
oblique section through the calcaneus and the head of the talus.



Fig. 1428 Foot, Pes;
sagittal section through the second toe.

→ 1315, 1316





Fig. 1429 Foot, Pes;
magnetic resonance tomography image (MRI); sagittal section.

→ 1317, 1318

Foot, cross-section and pedograms



Fig. 1430 Osteofibrous tubes of the foot; frontal section through the metatarsus.



Fig. 1431 a–c Foot prints, pedograms.
a Normal foot
b Pes cavus
c Flat foot

Index

The numbers refer to the page numbers.
The numbers with „T" refer to the numbers of the tables in the separate booklet enclosed in Volume 1.

A

Abdomen
- computed tomography 248–249, 251, 255
- cross-section 248–251
- frontal section 244–245
- median section 242
- sagittal section 243, 246–247

Abdominal viscera 158, 162, 166, 172
- arteries 168

Abdominal wall 58, 66, 69–71
- anterior, viewed from inside 59, 71
- innervation, segmental, sensory 67
- muscles 52, 54, 56–58, 60–61, 66, 72–73, T14–T16
- posterior 220–221
- skeleton projection 47
- surface anatomy 46

Acetabular floor 274
Acetabular roof 274, 280
Acetabulum 256, 261, 266
Achilles tendon s. Tendo calcaneus
Acini pancreatici 153
Acromion 1–2, 26, 47–48, 122
Adductor canal 357–358
Ala
- ossis ilium 72, 254, 264, 266–267, 273
- – sacri 10

ALCOCK's canal 230, 258
Ampulla
- ductus deferentis 71, 191, 195–196, 236
- (Duodenum) 130, 132, 134–135, 179
- hepatopancreatica 135, 154
- recti 210, 215, 242, 252–253, 257–258
- tubae uterinae 140, 167, 201, 203, 209, 253

Anal hiatus 230
Anastomoses, portocaval 181, 213
Angle
- between neck and shaft of the femur 276
- of antetorsion, femur 277

Angulus(-i)
- costae 49, 101
- inferior 1–2, 26–27
- infrasternalis 46–47
- mandibulae 12
- ossis pubis 266
- sterni 46, 50
- subpubicus 268
- superior 2

Ansa subclavia 117, 120
Anteflexio (Uterus) 203
Anteversio (Uterus) 203
Antrum pyloricum 130, 133, 135, 247
Anulus
- femoralis 229
- fibrosus 24–25, 29, 42
- – dexter (Cor) 77, 84
- – (Discus intervertebralis) 45
- – sinister (Cor) 77, 79, 84

Anulus
- inguinalis profundus 60, 71, 225
- – superficialis 56–58, 60–61, 70, 238–239
- umbilicalis 46, 52, 56, 58, 66, 70, 72–73, 140, 159, 242

Anus 210, 212–214, 230, 232–233, 235
Aorta 29, 35, 76–77, 83, 87, 122, 126, 174, 242, 249, 359
- abdominalis s. Pars abdominalis aortae
- ascendens s. Pars ascendens aortae
- congestion, collateral circulation 69
- descendens s. Pars descendens aortae
- thoracica s. Pars thoracica aortae

Apertura
- pelvis inferior 264–265
- – superior 264–265

Apex
- capitis fibulae 283, 291
- cordis 74–80, 84–85, 98–99, 101
- dentis 6
- ossis sacri 10–11
- patellae 284, 290–291
- pulmonis 94–95, 98–99, 110, 123
- vesicae 190–191, 252

Aponeurosis
- musculi obliqui externi abdominis 60–61
- plantaris 299, 304, 338, 340, 342, 376–379, 383–384

Appendix(-ces)
- epididymidis 193, 239
- epiploicae s. Appendix(-ces) omentales
- fibrosa hepatis 142–143, 160, 166
- omentales 138–139, 162, 165, 226, 258
- testis 192–193, 239
- vermiformis 138–140, 163–166, 175, 209, 251
- – projection 140
- – variations in position 140
- vesiculosae 200–201, 204

Arachnoidea mater
- cranialis 40
- spinalis 40–41, 43–45

Arcus
- anterior atlantis 6, 12, 19
- aortae 74–75, 84–85, 89, 98–99, 101, 104–105, 107–108, 114–115, 118–120, 122, 124
- – AP-radiograph 114
- – branches 114
- – variations 114
- cartilaginis cricoideae 90
- costalis 3, 46–47, 52, 207
- iliopectineus 71, 269, 313, 317, 355
- plantaris profundus 351, 375, 378
- posterior atlantis 6, 12, 18–19, 33, 39–40
- pubicus 268

Arcus
- tendineus musculi levatoris ani 230–231, 317
- – – solei 311, 329, 350, 370
- venae azygos 116
- venosus dorsalis pedis 365, 373
- – jugularis 110
- vertebrae 5–8, 15, 17, 20, 22, 29, 32, 41, 45

Area(-ae)
- cribrosa (Ren) 183
- gastricae 130
- intercondylaris anterior 282, 293
- – posterior 282
- nuda 119, 142, 145, 242

Areola mammae 64
Arteria(-ae)
- appendicularis 168, 174
- arcuata 183, 216, 351, 374
- ascendens 176
- auricularis posterior 38
- axillaris 54, 113, 120, 122, 124
- bulbi penis 234, 240
- – vestibuli 234
- caecalis posterior 174
- carotis communis 69, 74–75, 89, 98–99, 104–107, 110–111, 113–114, 117–118, 120–124
- – externa 114
- – interna 114
- caudae pancreatis 168
- cervicalis profunda 39, 115
- – superficialis 66
- circumflexa femoris lateralis 261, 351, 357–359
- – – medialis 260–261, 351, 357–359, 363–364
- – humeri posterior 37, 122
- – ilium profunda 179, 221, 227, 231, 356, 359
- – – superficialis 66, 353, 359
- – scapulae 37, 122
- colica dextra 152, 168, 174–175
- – media 152, 161, 163–164, 168, 174–177
- – sinistra 167–168, 175–177, 179, 217
- coronaria(-ae) 84, 86
- – dextra 74–75, 78–80, 82, 84–87, 98, 119–120, 127
- – – coronary angiography 87
- – sinistra 74–75, 79–80, 82, 84–87, 98, 119–120, 127
- – – coronary angiography 87
- cremasterica 193, 225, 238
- cystica 142, 149, 168, 170, 172
- descendens genus 351, 353, 356–358, 369
- digitales dorsales (Pes) 374
- – plantares communes 375, 377
- – – propriae 375–377
- dorsalis clitoridis 234
- – pedis 351, 368, 374–375, 379, 382
- – penis 199, 227, 234, 238–240, 244
- ductus deferentis 60, 193, 197, 227

Arteria(-ae)
- epigastrica inferior 60–61, 66, 69, 71–72, 158–159, 173, 175, 179, 196, 217, 221, 225, 227, 229, 231, 236, 253–254, 359
- – – collateral circulation 69
- – superficialis 66, 353, 359
- – superior 66, 69–70, 72
- – – collateral circulation 69
- femoralis 62, 69, 73, 218, 244, 256–257, 314, 317, 351, 353, 355–359, 380–381
- – superficialis 351, 356
- fibularis 351, 366, 368–372, 381–382
- gastrica(-ae) breves 131, 166, 168, 170, 173
- – dextra 131, 168, 170, 172
- – posterior 170
- – sinistra 106, 131, 152, 155, 160, 168, 170–173, 178–179, 218, 244, 247, 249
- gastroduodenalis 152, 168, 170–173, 242, 244
- gastroomentalis dextra 131, 167–168, 170–173
- – sinistra 131, 160, 168, 170–173, 247
- glutea inferior 212, 224–225, 256–257, 351, 363–364, 380
- – superior 212, 224–225, 254, 359, 363, 380
- helicinae 198
- hepatica communis 131, 152, 168, 170–173, 178–179, 218, 249
- – – AP-radiograph 171
- – propria 131, 142, 146, 152, 155, 167–168, 170–172, 179, 243–244, 249
- – variability 170
- ileales 152, 168, 174–175, 251
- ileocolica 168, 174–175, 255
- iliaca communis 167, 179, 212, 214, 217–218, 221, 224–229, 231, 242–243, 254, 359
- – externa 69, 71, 179, 196, 212, 221, 224–225, 227–229, 231, 236, 253, 317, 354, 356, 358–359
- – interna 212, 217, 224–225, 227–229, 231, 354, 356, 359
- – – variations 224
- iliolumbalis 217, 221, 224–225, 229
- inferior lateralis genus 289, 351, 368–369
- – medialis genus 289, 351, 358, 369–372
- intercostalis(-es) 70, 99, 120, 127
- – posteriores 45, 54, 73, 75, 104–105, 112–113, 115–116
- – suprema 115
- interlobares (Ren) 183–184, 216
- interlobulares (Hepar) 145
- jejunales 152, 168, 174–175, 177, 242, 244
- lienalis s. Arteria(-ae) splenica
- lobaris(-es) inferiores 98–99, 101, 126
- – media 98
- – superiores 98, 125

Arteria(-ae)
- lobi caudati 142
- lumbales 41, 179, 217, 221, 245, 359
- malleolaris anterior lateralis 351, 368, 374
- - - medialis 351, 374
- marginalis coli 168, 174–176
- media genus 289, 351, 369
- mesenterica inferior 152, 167–168, 174, 176–177, 179, 212, 214, 217, 221, 225, 227, 250–251
- - - AP-radiograph 176
- - - variations 176
- - superior 115, 152–154, 167–168, 170, 173–179, 187, 217–218, 221, 242–244, 250–251, 255
- - - AP-radiograph 174
- - - variations 174
- metatarsales dorsales 351, 368, 374
- - plantares 375–376, 378
- musculophrenica 69
- obturatoria 71, 212, 224–225, 227, 229, 231, 236–237, 256–257, 261, 317, 355, 357–359
- - variability 229
- occipitalis 37–39
- ovarica 152, 167, 201, 204, 209, 223, 228–229, 253
- pancreatica dorsalis 168, 171
- - inferior 171
- - magna 168
- pancreaticoduodenalis inferior 168, 173, 177
- - superior anterior 171, 173
- - - posterior 168, 170–171, 178
- perforantes 351, 358–359, 362–363, 369
- pericardiacophrenica 69, 89, 112–113, 118
- perinealis 234, 240
- phrenica inferior 63, 104, 172–173, 189, 217–218, 221
- plantaris lateralis 304, 351, 375, 377–378, 382
- - medialis 304, 351, 375, 378, 382
- - profunda 374–375
- poplitea 284, 289, 320, 329, 351, 358, 361–363, 366, 369–372
- - variations 370
- prepancreatica 168
- profunda brachii 37
- - femoris 256–257, 261, 351, 356–359, 380
- - penis 198–199, 226–227, 239, 244
- pudenda externa 66, 238, 353, 359
- - interna 210, 212, 224–225, 227–229, 231, 234, 236–237, 240–241, 256–258, 260–261, 363–364
- pulmonalis(-es) 112
- - dextra 74–76, 84–85, 88–89, 95, 98–101, 105, 107, 119, 125
- - sinistra 74–75, 84–85, 88–89, 95, 98–99, 105, 107, 113, 124
- rectalis 212
- - inferior 168, 212, 227–228, 240–241
- - media 212, 224–225, 227–229
- - superior 167–168, 176–177, 179, 210, 212, 217, 227–229, 254

Arteria(-ae)
- recurrens tibialis anterior 351, 368
- - - posterior 351, 369–370
- renalis 152, 182–183, 185, 187, 189, 216–218, 221, 242–243, 245–246, 250
- - accessoria 217
- - arteriography 216
- - polaris inferior 216
- - - superior 216
- - variations 216
- sacralis lateralis 224–226
- - mediana 177, 179, 212, 217, 221, 225, 227–229, 253, 359
- segmentalis basalis anterior 98
- - - lateralis 99
- - - medialis 99
- - - posterior 99
- segmenti (A. renalis) 216
- - anterioris hepatici 124
- - - renalis 171
- - posterioris hepatici 124
- - - renalis 171
- sigmoideae 167–168, 176–177, 179, 212, 217, 254, 258
- spinalis anterior 45
- splenica 152, 154–156, 161, 167–168, 170–173, 178–179, 218, 245, 248–249
- subclavia 74–75, 89, 98–99, 104–107, 111–113, 115, 117, 123
- - dextra 69, 99, 114, 118, 120
- - sinistra 99, 114, 120, 124
- subcostalis 115–116, 245
- subscapularis 124
- superior lateralis genus 351, 366, 368–369
- - medialis genus 351, 358, 362, 368–369
- supraduodenalis 168
- suprarenalis 189
- - inferior 182, 189, 217
- - media 182, 189, 217
- - superior 182, 189
- - variations 189
- suprascapularis 120
- surales 351, 369–370
- tarsalis(-es) lateralis 351, 374
- - mediales 351, 374
- testicularis 70–71, 152, 179, 193–194, 197, 217, 221–222, 238–239, 243, 251, 254, 256
- - variations 217
- thoracica interna 63, 66, 69, 88, 110, 112, 114–115, 118, 120, 124, 126
- - - collateral circulation 69
- - lateralis 66
- thoracoacromialis 120
- thoracodorsalis 124
- thyroidea inferior 106
- - superior 110
- tibialis anterior 283, 351, 366–370, 372, 374, 381
- - posterior 351, 366, 369–372, 375–378, 381–382
- transversa colli 38–39
- umbilicalis 70–71, 159, 185, 207, 224–225, 227–229, 236
- urethralis 199
- uterina 204, 224, 228–229
- - Variabilität 204
- - variability 204
- vaginalis 204, 228–229
- vertebralis 24, 34, 37, 39–40, 106, 114, 120

Arteria(-ae)
- vesicalis(-es) inferior 225, 227–228
- - superiores 70, 224, 227, 229, 260

Arteries, hip 359
Arteriography 216
Arthroscopy, Articulatio genus 292
Articulatio(-nes)
- acromioclavicularis 122
- atlantoaxialis lateralis 17–19
- - mediana anterior 19
- - - posterior 19
- atlantooccipitalis 17–19
- calcaneocuboidea 263, T39
- capitis costae 8, 14, 20, 22
- costotransversaria 8, 14, 20
- costovertebrales 20–22
- coxae 257, 263, 268–271, 273, 278–279, T39
- - AP-radiograph 280
- - LAUENSTEIN projection 280
- cuneocuboidea 263, T39
- cuneonavicularis 263, T39
- femoropatellaris 263, 284
- genus 263, 284–289, 292, 323, T39
- - AP-radiograph 291
- - arthroscopy 292
- - cross-section 284
- - frontal section 290
- - lateral radiograph 291
- - MRI 293
- - sagittal section 290
- humeri 122–123, T39
- intercuneiformes 263, T39
- intermetatarsales 304, T39
- interphalangeae pedis 263, T39
- lumbosacralis 268–271
- membri inferioris 263, T39
- - - liberi 263, T39
- meniscofemoralis 263
- meniscotibialis 263
- metatarsophalangea(-ae) 263, 302, 304, T39
- - hallucis 336
- pedis 300–303
- sacrococcygea 226, 252, 264
- sacroiliaca 1, 16, 176, 205, 230, 254, 263–264, 268–269, 271–273, 280, 317, T39
- sternoclavicularis 46, 51, 119
- sternocostales 51
- subtalaris 263, 304–305, 383, T39
- talocalcaneonavicularis 263, 303–305, T39
- - AP-radiograph 305
- - frontal section 304
- - lateral radiograph 305
- - sagittal section 304
- talocruralis 263, 301, 304–305, 383, T39
- - AP-radiograph 305
- - frontal section 304
- - lateral radiograph 305
- - sagittal section 304
- talonavicularis 383, T39
- talotarsalis 263, 304, T39
- tarsi transversa 296, T39
- tarsometatarsales 263, 296, 304, T39
- tibiofibularis 263, 282, 290–291, 293–294, T39
- zygapophysialis 12, 15–16, 23, 25, 29, 35, 272

Atlas 4–7, 12, 17–19, 31, 33–34, 39–40
Atrium
- cordis dextrum 74–76, 78, 82, 84–85, 89, 99, 101, 114, 119–121, 127, 243–244, 255
- - sinistrum 75–76, 79, 82, 84–85, 89, 99, 101, 114, 119, 126–127

Auricula
- dextra 74, 78–80, 84, 98, 114
- sinistra 74–76, 79, 84–85, 98, 101, 114, 126

Axilla 122–123
Axis 4, 6–7, 12, 17–19, 33
- pelvis 265

B

Back 1, 37
- cutaneous innervation, segmental 36
- dermatomes 36
- lines of orientation 1
- muscles 26–34
- regions 1
- surface anatomy 1

BARTHOLIN's gland 200, 233, 235
Basis
- ossis metatarsi 295, 298, 302
- - - hallucis 296
- - sacri 10–11, 15, 273
- patellae 284, 290–291
- phalangis (Pes) 295
- prostatae 195
- pulmonis 95
- pyramidis 183

Bifurcatio
- aortae 177, 179, 359
- tracheae 90–91, 99, 102, 104–105, 107, 119, 125
- trunci pulmonalis 75–76, 85, 89

BOCHDALEK's triangle 63
BOYD's vein 352
Bronchi, segmental s. Bronchus(-i) segmentalis(-es)
Bronchiolus(-i) 98
- terminalis 98

Bronchography/Bronchoscopy 93
- Bifurcatio tracheae 102
- segmental bronchi 102

Bronchus(-i) 90–92, 98
- arterial supply 99
- bronchography 93
- bronchoscopy 102
- intrasegmentalis 120, 124
- lingularis inferior (B V) 90, 93, 102
- - superior (B IV) 90, 93, 102
- lobaris(-es) 92, 107
- - inferior 126
- - - dexter 90–91, 95, 120–121
- - - sinister 90–91, 93, 98
- - medius 90–91, 95, 121, 126
- - superior 125
- - - dexter 90–91, 95, 99, 104, 125
- - - sinister 90–91, 93, 98–99, 102
- principalis dexter 88, 90–91, 95, 99, 101–102, 104–105, 107, 109, 112, 115, 117, 119–120, 125
- - sinister 90–91, 93, 95, 99, 102, 104–105, 113, 115, 117, 119, 125
- projection 90

Bronchus[-i]
- segmentalis[-es] 90–92, 99
- - anterior [B III] 90, 93, 102, 122, 124–125
- - apicalis [B I] 90
- - apicoposterior [B I + II] 90, 93, 102
- - basalis anterior [B VIII] 90, 93, 102
- - - cardiacus [B VII] s. Bronchus[-i] segmentalis basalis medialis [B VII]
- - - lateralis [B IX] 90, 93, 102, 127
- - - medialis [B VII] 90
- - - posterior [B X] 90, 93, 102, 126
- - bronchoscopy 102
- - lateralis [B IV] 90
- - medialis [B V] 90
- - posterior [B II] 90, 124–125
- - superior [B VI] 90, 93, 102

BRUNNER's glands 136
Bulbus
- aortae 79, 87
- penis 196, 198, 236, 252
- vestibuli 200, 228, 235, 237, 241, 261

Bursa[-ae]
- anserina 316
- iliopectinea 310, 355
- infrapatellaris profunda 285, 288–289
- ischiadica musculi obturatorii interni 311
- musculi semimembranosi 329–331
- omentalis 157, 161, 166–167, 247–249
- subcutanea calcanea 338
- - coccygea 37
- - infrapatellaris 306, 313
- - prepatellaris 284, 306
- - sacralis 37
- - spinae iliacae posterioris superioris 37
- - trochanterica 311
- subfascialis prepatellaris 288
- subtendinea iliaca 256, 315–316, 380
- - musculi gastrocnemii medialis 311, 329–331
- - - obturatorii interni 256–257
- - - sartorii 316
- - prepatellaris 313
- suprapatellaris 285, 288, 290, 292
- trochanterica musculi glutei maximi 279, 319–321
- - - - medii 256, 279, 311, 321

C

Caecum 138–140, 158, 162–166, 175, 209, 251, 254
- projection 140

Calcaneus 295–301, 303–304, 383
Calices renales 216
- majores 183, 186, 245
- minores 183–184, 186, 246, 250

Calx 262
Canal of ALCOCK 230, 258
Canalis[-es]
- adductorius 314, 351
- analis 211, 215, 252
- cervicis uteri 201–203, 253

Canalis[-es]
- inguinalis 239
- nervi hypoglossi 18
- obturatorius 212–213, 230, 268, 278–279, 316–317, 355
- pudendalis 230
- pyloricus 130, 135, 152
- sacralis 10–11, 24, 227, 252, 271, 317
- vertebralis 3, 24, 41–43, 45, 317
- - veins 42

Capsula
- adiposa 182, 188, 217, 245–247, 249
- articularis (Articulatio atlantoaxialis lateralis) 17–18
- - (Articulatio atlantooccipitalis) 18
- - (Articulatio costovertebralis) 21–22
- - (Articulatio coxae) 268, 270–271
- - (Articulatio genus) 285, 290, 293, 323
- - (Articulatio metatarsophalangea hallucis) 336
- - (Articulatio sternoclavicularis) 51
- - (Articulatio zygapophysialis) 25
- - (Columna vertebralis) 42
- fibrosa (Ren) 182–184, 246–247

Caput
- breve (M. biceps brachii) 54
- - (M. biceps femoris) 318, 320–323, 363, 381, T40, T46
- costae 8, 14, 22, 49
- epididymidis 192–194, 239, 252
- femoris 256, 259–261, 271, 273, 275–277, 279–280, 359
- fibulae 262, 282–283, 285–287, 291, 293–294, 306, 313, 322, 332, 367
- humeri 122–123
- laterale (M. gastrocnemius) 284, 286–288, 320–323, 326, 328–330, 362, 369–370, T49
- - (M. triceps brachii) 52
- longum (M. biceps brachii) 122
- - (M. biceps femoris) 317–323, 362–363, 380–381, T46
- - (M. triceps brachii) 37, 52
- mediale (M. gastrocnemius) 284, 286–287, 308, 320–321, 323, 326, 328–331, 362, 369–370, 381, T49
- obliquum (M. adductor hallucis) 304, 339, 342, 378, 383–384, T52
- ossis metatarsi 295, 298
- - - quinti 296
- pancreatis 134, 140, 152–153, 167, 173, 179, 243–244
- phalangis (Pes) 295
- rectum 278–279, 315, T44
- reflexum 278–279, 315, T44
- tali 295–298, 305, 382–383
- transversum (M. adductor hallucis) 339–342, 378, T40, T52

Cardia 109, 130–131, 160, 242, 247
Carina tracheae 102, 119
Cartilago[-ines]
- articularis 284
- arytenoidea 91
- bronchiales 90, 102
- corniculata 91
- costalis 51, 55, 57–58, 70, 115, 125, 127

Cartilago[-ines]
- cricoidea 12, 119
- epiphysialis 274
- fibrosa (Discus intervertebralis) 272
- hyalina (Discus intervertebralis) 272
- thyroidea 35, 58, 90–91, 104, 110, 119–120
- tracheales 90–91, 104–105, 119

Caruncula hymenales 233, 235
Cauda
- epididymidis 192–194, 252
- equina 24, 29, 41, 43–44, 242, 250–251, 254
- pancreatis 134, 152, 154–155, 161, 166–167, 179, 244, 249

Cave of RETZIUS 207, 252
Cavea
- thoracis 48, 55
- - AP-radiograph 101

Caverna[-ae]
- corporis cavernosi 198–199
- - spongiosi 199

Cavitas[-tes]
- articularis (Articulatio coxae) 271, 273, 279–280
- glenoidalis 48, 122
- infraglottica 119
- medullaris 277, 281
- nasi 12
- pericardiaca 119, 125
- peritonealis 157, 167, 242–244, 247–251
- - development 157
- pleuralis 112–113, 117, 121, 126, 155, 248–249
- serosa scroti 61
- symphysialis 272
- thoracis 119, 124–125, 127
- - CT 129
- - frontal section 120–122
- - median sagittal section 119
- - MRI 120–121
- - transverse section 123, 125–126
- uteri 201–203, 237, 253, 259

Cavum serosum scroti 61, 194
Centrum
- perinei 230–231
- tendineum 55, 62–63, 73, 104, 106, 112–113, 242, 246

Cerebellum 40
Cerebrospinal fluid, puncture 44
Cervical part of the vertebral column 17, 43
- AP-radiograph 13
- CT 35
- lateral radiograph 12

Cervix
- uteri 203–205, 237
- vesicae 195

Chiasma
- crurale 326
- plantare 342
- tarsale 326

CHOPART's joint 296, T39
Chorda[-ae]
- arteriae umbilicalis 61, 66, 71–72, 158, 196, 229, 236, 253
- tendineae 79–81
- urachi 61, 72, 236

Cingulum
- membri inferioris s. Cingulum pelvicum
- - superioris s. Cingulum pectorale

Cingulum
- pectorale 3, 48
- pelvicum 3, 263–264

Cisterna
- cerebellomedullaris 40
- chyli 116, 218

Clavicula 3, 46–48, 51–52, 90, 101, 108, 110, 112–113, 118, 120, 122–123, 159
Cleft, uncovertebral 24
Clitoris 207, 223, 231
Clivus 18
COCKETT's veins 352
Coeliacography 171
Colliculus seminalis 190, 198
Collum 257
- costae 8, 20, 49
- femoris 259, 273, 275–280
- fibulae 283
- tali 296–298, 305
- vesicae biliaris 149–151

Colon 138–139, 188
- AP-radiograph 141
- ascendens 138–141, 158, 161–162, 165–166, 180, 186, 244, 251, 254
- descendens 138, 140, 159, 165–166, 177, 180, 186, 244–245, 249–251, 254–255
- layers of the wall 139
- projection 138
- sigmoideum 138, 140, 157–158, 162, 164–167, 175, 177, 179, 207, 209, 212–213, 215, 217–218, 226, 242, 244, 251–254, 258
- transversum 138–140, 149, 157–158, 160–162, 164–166, 175, 242–244, 246–247, 249–250
- - variations in position 141

Colonoscopy 141, 211
Columna[-ae]
- anales 210–211, 242, 252, 258
- renales 183, 246
- rugarum anterior 235
- vertebralis 2–4, 42
- - nerves 42

Commissura
- labiorum anterior 235
- - posterior 235

Compartimentum
- cruris anterius 366
- - laterale 366
- - posterius 366

Complexus stimulans cordis 82–83
Computed tomography (CT)
- abdomen 249, 251
- cervical part of the vertebral column 35
- lumbar part of the vertebral column 35
- Mm. abdominis 73
- Mm. dorsi 29
- pelvis 254
- - of the female 257
- - of the male 256
- thoracic cavity 125

Conducting system 82–83
Condylus
- lateralis (Femur) 275, 277, 281, 285–287, 290–293
- - (Tibia) 282, 287, 290–293
- medialis (Femur) 275, 277, 281, 285–287, 290–293, 313–314
- - (Tibia) 282, 290–291, 293, 327, 329, 331–332
- occipitalis 6

Conjugata
- anatomica 265
- diagonalis 265
- vera 264–265
Conus arteriosus 74, 84, 98
Cor 67, 74–75, 98, 114, 242
- arterial supply 85
- atrium, left 79
- - right 78, 82
- balanced coronary artery supply 85–86
- conducting system 82–83
- coronary arteries 84, 86–87
- - coronary angiography 87
- dominant left coronary artery supply 85–86
- dominant right coronary artery supply 85–86
- innervation 83
- position 77
- - during exspiration/inspiration 88
- projection 101
- ultrasound image 126
- valves, points of auscultation 77
- - projection 77
- ventricle, left 79–82
- - right 78, 80–82
Cornu
- anterius (Meniscus lateralis) 290, 293
- coccygeum 11
- inferius (Margo falciformis) 306, 355
- posterius (Meniscus lateralis) 290, 293
- sacrale 10–11
- superius (Margo falciformis) 306, 355
Corona glandis 196, 198–199, 252
Coronary angiography 87
Coronary arteries 84, 86
Corpus(-ora)
- adiposum infrapatellare 285, 290, 293, 323
- - pararenale 188
- - (Ren) 183
- albicans 201
- anococcygeum 207, 226, 230–233, 235, 240–241, 252, 257
- cavernosum clitoridis 207, 235, 253
- - penis 192, 196–197, 199, 222, 227, 239, 242–243, 252, 259
- - recti 211
- (Clavicula) 46–47
- clitoridis 231, 253
- costae 49, 101
- epididymidis 191, 194
- femoris 215, 275, 277, 291
- fibulae 283, 291
- gastricum 121, 128, 130, 132–133, 158, 160–161, 245
- luteum 201–202
- - menstruationis 202
- mammae 52
- ossis ilium 259, 267
- - ischii 266–267
- - metatarsi 295, 298, 334
- - pubis 230, 243, 266–267
- pancreatis 134, 152–155, 161, 166, 173
- penis 197
- phalangis (Pes) 295
- spongiosum penis 192, 196–199, 222, 232, 239, 242, 252, 259–260
Corpus(-ora)
- sterni 3, 47, 50–51, 55, 90, 119, 124–126, 242
- tali 296–297, 304
- tibiae 282–283, 291
- uteri 201, 203–204, 237, 253, 259
- vertebrae 5–9, 12–25, 29, 35, 41, 45, 72, 125, 155, 188, 249, 251
- vesicae 190–191
- - biliaris 149–151
- - cystoscopy 190
Corpusculum renale 184
Cortex
- (Glandula suprarenalis) 189
- renalis 183–184, 188, 246–247, 250
Costa(-ae) 2–3, 5, 8, 14–16, 20–22, 27, 29, 31–32, 34, 41, 48–49, 54, 58, 63, 69, 73, 103, 110, 113, 115–116, 118–120, 122, 124–127, 132, 144, 183, 188, 216–217, 220, 245–250
- fluctuantes (XI; XII) 48
- spuriae (VIII-XII) 48
- verae (I-VII) 48
COWPER's gland 226, 232, 234
Coxa
- valga 276
- vara 276
Craniocervical junctions 17–19
- AP-radiograph 19
Crista
- capitis costae 49
- colli costae 49
- iliaca 1–2, 15, 26–27, 37, 44, 47, 62, 73, 103, 176, 183, 216, 220, 230, 254, 264, 266–267, 271, 273, 307, 317, 319–320, 322, 360, 364
- intertrochanterica 273, 275–276
- medialis 283
- obturatoria 266–267
- occipitalis externa 6
- pubica 266
- sacralis lateralis 10
- - medialis 10
- - mediana 10–11, 15, 360
- urethralis 190, 198
Crus(-ra) 263
- clitoridis 200, 235, 237, 261
- compartments 366
- cross-section 381–382
- cutaneous innervation 350
- dextrum (Fasciculus atrioventricularis) 82–83
- - (Pars lumbalis, Diaphragma) 62–63, 68, 104, 106, 116, 155, 249, T21
- extensors 326
- flexors 326
- laterale (Anulus inguinalis superficialis) 56, 58, 60, 239
- mediale (Anulus inguinalis superficialis) 56, 58, 60, 239
- muscles 324–332, T47–T50
- - lateral 366
- - origins and insertions 310–311, 324–325
- osteofibrous tubes 366
- penis 196, 198, 226, 236, 260
- sinistrum (Fasciculus atrioventricularis) 82–83
- - (Pars lumbalis, Diaphragma) 63, 104, 106, 116, 248, T21
Cryptae intestinales s. Glandulae intestinales
Cumulus oophorus 202
Cupula pleurae 103, 122
Curvatura
- major 128–130, 132, 158–161
- minor 128–130, 132, 135, 160–161
Cuspis
- anterior (Valva atrioventricularis dextra) 77–78, 81–82
- - (Valva atrioventricularis sinistra) 77, 79–80, 82, 126
- commissuralis dextra 77
- - sinistra 77
- posterior (Valva atrioventricularis dextra) 77–78, 80–81
- - (Valva atrioventricularis sinistra) 77, 79–81, 126
- septalis (Valva atrioventricularis dextra) 77–78, 80
Cutaneous innervation, segmental
- back 36
- lower limb 346
Cutaneous nerves, lower limb 349
Cutis 70, 188, 210
Cystoscopy 190–191

D

Defaecography 215
Dens axis 6–7, 12, 18–19
Dermatomes
- back 36
- lower limb 346
Diameter(s)
- obliqua I 264
- obliqua II 264
Diameter(s)
- of the pelvis in the female 265
- sagittalis 265
- transversa 264–265
- tuberalis 265
Diaphragma 54–55, 59, 62–63, 67, 69, 73, 87, 101, 104–112, 116–121, 142, 144, 148–149, 152, 157, 159–160, 255, T21
- pelvis 188, 237, 242, 245–246, 249–250, T22
- - of the female 230–231, 233
- - of the male 232, 236
- urogenitale 237, 241, 243, 261, T22
- - of the female 234–235
- - of the male 234
Digitus(-i)
- minimus (Pes) 262, 295
- pedis 263
- primus (Pes) s. Hallux
- quartus (Pes) 262, 295
- quintus (Pes) s. Digitus minimus
- secundus (Pes) 262, 295
- tertius (Pes) 262, 295
Discus
- articularis (Articulatio sternoclavicularis) 51
- interpubicus 207, 268, 271–272, T39
- intervertebralis 4, 12, 17, 19–25, 29, 35, 42, 45, 178, 259, 268, 271
Distantia
- intercristalis 265
- interspinosa anterior 265
- - posterior 265
- pubococcygea 265
Diverticulum(-a)
- ampullae 191
- cervical 109
- epiphrenic 109
- ilei 137
- mid-oesophageal 109
- of MECKEL 137
- of ZENKER 109
DODD's veins 306, 352
Dorsum
- lines of orientation 1
- of the foot, muscles 333–337, T51–T52
- pedis 262, 368, 373–374
- penis 197
- regions 1
- surface anatomy 1
DOUGLAS, Pouch of 157, 167, 207, 229, 253
Duct
- of MÜLLER 192, 200
- of SANTORINI 152
- of WIRSUNG 152
- of WOLFF 192, 200
Ductulus(-i)
- aberrans inferior 192, 194
- efferentes testis 194, 252
- prostatici 190, 198
- transversi 200–201
Ductus(-us)
- alveolares 98
- arteriosus [BOTALLO] 75
- biliaris s. Ductus choledochus
- bilifer interlobularis 145
- choledochus 142, 146, 150–154, 170, 172, 179, 243, 249
- - AP-radiograph 151, 154
- - variability 150, 153
- cysticus 131, 150–154, 167, 170, 179, 249
- deferens (Vas deferens) 60–61, 70–71, 166, 179, 190–192, 194–197, 221–222, 226–227, 236, 238–239, 243, 252, 256, 258
- - AP-radiograph 195
- ejaculatorius 190, 192, 195–196, 198
- epoophorontis longitudinalis s. Ductus longitudinalis
- excretorius 153, 196
- glandulae bulbourethralis 192, 198, 234
- hepaticus communis 131, 150–154, 167, 170, 179
- - dexter 146, 150–151
- - sinister 150–151
- lactiferi 64
- longitudinalis 200–201
- mesonephricus 192, 200
- omphaloentericus 137
- pancreaticus 150, 152–154, 242–243
- - accessorius 152–153
- - AP-radiograph 154
- - variability 153
- paramesonephricus 192, 200
- paraurethralis 235
- thoracicus 63, 113, 115–116, 121, 124–127, 245, 248–249
- venosus [ARANTIUS] 143
Duodenum 117, 128–130, 133–136, 140, 149–154, 157, 160–161, 163, 165–167, 170, 179–180, 243–244, 255
- AP-radiograph 132, 135
- endoscopy 135
- layers of the wall 136

Duodenum
- projection 134
Dura mater
- cranialis 19
- spinalis 19, 40–41, 43–45, 119

E

Embryo, ultrasound image 206
Eminentia
- iliopubica 230, 264, 266–267, 273
- intercondylaris 282, 290–291
- plantaris intermedia 384
- - lateralis 338, 384
- - medialis 338, 384
Endocardium 80
Endoscopic retrograde cholangiopancreaticography (ERCP) 154
Epicardium s. Lamina visceralis (Pericardium serosum)
Epicondylus
- lateralis (Femur) 275, 281, 284, 290–291
- medialis (Femur) 275, 277, 281, 284, 287, 290–291, 315
Epididymis 61, 185, 192–194, 222, 239, 242–243
Epiglottis 12
Epineurium 43
Epiorchium 194
Epiphysis
- anularis 5, 7–9, 24–25
- of the head of the femur 274
Episiotomy 233
Epithelium 105, 130, 136, 139
Epoophoron 200
Erection 222
Excavatio
- rectouterina 157, 167, 201–202, 205, 207, 229, 253, 257
- rectovesicalis 166, 190, 196, 210, 217, 226, 236, 243, 252
- vesicouterina 157, 167, 202, 205, 207, 209, 237, 253
Extremitas
- acromialis 47, 122
- anterior (Splen) 156, 159
- inferior (Ren) 164, 167, 186, 188, 216
- - (Testis) 193
- posterior (Splen) 156
- sternalis 3, 47, 108
- superior (Ren) 160, 166, 188, 216
- - (Testis) 193
- tubaria 201
- uterina 201

F

Facies
- anterior (Glandula suprarenalis) 189
- - (Patella) 284–285, 290–291
- - (Ren) 188
- anteroinferior 164
- anterosuperior 161
- articularis anterior (Axis) 6
- - calcanea 298
- - - anterior 297
- - - media 297
- - - posterior 297
- - capitis costae 49
- - - fibulae 283

Facies articularis
- - cuboidea 297–298
- - fibularis 282
- - inferior (Atlas) 6
- - - (Tibia) 282, 294, 301
- - - (Vertebra lumbalis) 9
- - malleoli lateralis 283, 294, 301
- - - medialis 282, 294, 301
- - navicularis 297–298
- - (Patella) 284–285, 290, 292
- - posterior (Axis) 6
- - superior (Atlas) 6, 19
- - - (Tibia) 282, 293
- - talaris 303
- - - anterior 297–298, 303
- - - media 297–298, 303
- - - posterior 297–298, 303
- - tuberculi costae 49
- auricularis 10–11, 266
- colica 156
- contacta hepatis 217
- costalis (Pulmo) 94–95
- diaphragmatica (Hepar) 142, 145, 160–161
- - (Pulmo) 95
- - (Splen) 156
- dorsalis (Os sacrum) 26
- gastrica (Splen) 156, 166–167
- glutea 267
- intervertebralis 7–9, 12, 14–16, 25
- intestinalis 201–202, 209, 229, 253
- lateralis (Fibula) 283
- - (Testis) 193
- - (Tibia) 282–283
- lunata 230, 266–267, 279
- malleolaris lateralis 296–298
- - medialis 296–298
- medialis (Fibula) 283
- - (Ovarium) 201
- - (Tibia) 282–283, 306, 327, 367
- patellaris 275, 281, 284–286, 290, 292
- pelvica 11
- poplitea 275, 277, 291, 323, 328, 330, 369
- posterior (Fibula) 283
- - (Prostata) 191
- - (Ren) 188
- - (Tibia) 282–283
- renalis (Splen) 156
- sacropelvica 266
- superior (Talus) 297
- symphysialis 3, 266, 271, 279
- vesicalis 202, 209, 253
- visceralis (Hepar) 145, 160
- - (Splen) 156
Falx septi 79
Fascia
- axillaris 58
- brachii 58
- cervicalis 110, 119
- clavipectoralis 58
- cremasterica 61, 193–194, 197, 239, 252
- cribrosa 218, 306
- cruris 306–307, 328, 365–366, 381–382
- deltoidea 26–27
- dorsalis pedis 306, 379, 384
- glutea 52, 307, 319, 362
- iliaca 179
- inferior diaphragmatis pelvis 230
- infraspinata 26–28

Fascia
- investiens perinei superficialis s. Fascia perinei
- lata 306–307, 313–315, 319, 355, 360, 381
- nuchae 39
- obturatoria 230, 232–233, 317
- pedis 382
- pelvis parietalis 253
- - visceralis 237, 252
- penis profunda 197, 199, 238
- - superficialis 199
- perinei 230, 232–235, 260
- rectoprostatica 252
- rectovaginalis 207, 253
- renalis 188, 246–247
- spermatica externa 60–61, 70, 197, 239
- - interna 61, 70, 193–194, 197, 239
- superficialis (Canalis inguinalis) 61
- thoracolumbalis 22, 24–32, 35, 52, 72, 319
- transversalis 31, 58–62, 69, 72–73
- vesicovaginalis 207
Fasciculus(-i)
- atrioventricularis 82–84
- lateralis 122
- longitudinales (Lig. cruciforme atlantis) 18–19
- medialis 122
- posterior 122
- transversi (Aponeurosis plantaris) 338
Fat pad, parathymic 111
Femur 3, 215, 256–257, 262–263, 273–277, 280–281, 284–287, 290–293, 313–316, 323, 328, 330, 369, 381
- angle between neck and shaft of the femur 276
- - of antetorsion 277
- AP-radiograph 274
- arteries 351, 359
- - digital subtraction angiography (DSA) 359
- cross-section 380–381
- fascia 306–307
- muscles 310–323, T42–T46
- - origins and insertions 310–311
- structure of the trabecular bone 276
- veins 352
Fibrae
- intercrurales 56, 58, 60
- obliquae 129
Fibula 262–263, 282–283, 285–287, 294, 300–301, 304–305, 330–331, 335, 366, 381–382
Filum(-a)
- radicularia (N. spinalis) 45
- - posteriora 40
- terminale 43, 242, 252
Fimbria(-ae)
- ovarica 201
- tubae uterinae 201, 209, 253
Fissura(-ae)
- horizontalis pulmonis dextri 94–95, 120
- ligamenti teretis 142
- obliqua 94–95, 99, 120–121, 124–127

Fissura(-ae)
- portalis dextra 147
- - principalis 147
- umbilicalis 147
Flatfoot 303, 384
Flexio (Uterus) 203
Flexura
- anorectalis 210, 226
- coli dextra 138, 155, 160–163, 165–166, 175
- - sinistra 120–121, 132, 138, 162, 165–166, 175
- duodeni inferior 131, 153, 175
- - superior 153
- duodenojejunalis 134–135, 152, 166–167, 177, 244
- perinealis s. Flexura anorectalis
- sacralis 210, 226
Foetus, ultrasound image 206
Fold of KOHLRAUSCH 210, 252
Folliculus(-i)
- atreticus 202
- ovaricus(-i) maturus 202
- - vesiculosi 201, 253
- primarii 202
Foot s. Pes
Foot prints 384
Foramen(-ina)
- costotransversarium 8, 21
- epiploicum s. Foramen omentale
- infrapiriforme 317, 319–321, 360, 364
- intervertebrale 4, 12, 14–15, 22, 25, 29, 35, 41, 45, 271
- ischiadicum majus 230, 265, 269–271, 319, 360
- - minus 265, 269–271, 317, 319, 321
- magnum 6, 40
- nutricium 282–283, 311
- obturatum 229, 265–266, 269, 271, 273–274, 280
- omentale 160, 166–167
- ovale (Cor) 78
- papillaria 183
- sacralia anteriora 10, 16, 317
- - posteriora 10–11
- suprapiriforme 317, 319–320, 360
- transversarium 6–7, 35
- venae basivertebralis 9, 25
- - cavae 55, 62–63, 104, 106, T21
- venarum minimarum 78
- vertebrale 6–9, 35, 125
- WINSLOWI 160
Fornix
- gastricus 130, 132, 242, 247
- vaginae 202, 205, 207, 253
Fossa
- acetabuli 71, 230, 256, 260, 266–267, 273, 279–280
- iliaca 264, 266, 271
- inguinalis lateralis 59, 61, 71, 236
- - medialis 59, 61, 71, 236
- intercondylaris 275, 277, 281, 291, 293
- ischioanalis 232–233, 236, 256–258
- malleoli lateralis 283
- navicularis urethrae 198–199, 252
- ovalis 78
- paravesicalis 59, 260–261
- poplitea 262, 307, 321, 360–363, 366, 369–372

Fossa
- supravesicalis 61, 71, 236
- trochanterica 275–277, 279
- vestibuli vaginae 235
Fovea(-ae)
- capitis femoris 273, 275–277, 279–280
- costalis inferior 8, 20
- - processus transversi 8–9, 21, 45
- - superior 8–9, 20–21
- dentis 6, 19
Foveolae gastricae 130
FRANKENHÄUSER's ganglion 223
Frenulum
- clitoridis 235, 253
- labiorum pudendi 235
- ostii ilealis 139
- preputii 199
Fundus
- gastricus 120, 128–130, 132, 149, 160
- uteri 167, 201–204, 207, 209, 253
- vesicae 167, 195–196, 198
- - biliaris 149–150, 158, 163, 166
Funiculus
- spermaticus 56–58, 60, 68, 193, 197, 218, 226–227, 238, 256, 306, 354–355
- umbilicalis 70, 206, 208

G

Ganglion(-ia)
- cardiacum 120
- cervicale medium 83, 117, 120
- - superius 83
- cervicothoracicum 83, 117, 120
- coeliaca 214, 249
- of FRANKENHÄUSER 223
- mesentericum inferius 214
- pelvica 214, 222–223
- sacralia 225
- sensoria nervorum spinalium 24, 40–43, 45
- stellatum s. Ganglion cervicothoracicum
- thoracica 113–114, 117, 120–121
- trunci sympathici 41–43, 45, 214, 222–223
Gaster 63, 67, 104, 106, 108–109, 117, 121, 128–135, 140, 155, 157–161, 166, 170, 172, 179–180, 217, 242–244, 246–250, 255
- AP-radiograph 132, 171
- arteries, variability 170
- forms 132
- layers of the wall 130
- projection 132
Gastroscopy 133
Genital organs
- female 223
- - development 200
- - external 233, 235, 241
- - innervation 223
- - internal 204, 209
- male 222, 239
- - development 192
- - external 197, 238, 240
- - innervation 222
Genu 263
- muscles 323
GIMBERNAT's ligament 355
Gland(s)
- of BARTHOLIN 200, 233, 235
- of BRUNNER 136
- of COWPER 226, 232, 234
Glandula(-ae)
- areolares 64
- bulbourethralis 192, 196, 198, 222, 226, 232, 234, 236
- duodenales 136
- gastricae 130
- intestinales 136, 139
- mammaria 125–127
- oesophageae 105
- suprarenalis 152, 161, 167, 179, 182, 185, 189, 217, 245, 249
- thyroidea 104–105, 107, 110–111, 118–119
- tracheales 91
- vesiculosa 191–192, 195–196, 222, 226–227, 236, 242–243, 256, 258
- - AP-radiograph 195
- vestibularis major 200, 233, 235, 237
Glans
- clitoridis 200, 207, 233, 235
- penis 192, 196–199, 222, 242–243, 252
GLISSON, Triad of 145
Glomus coccygeum 253
GRAAFian follicle 201
Gubernaculum testis 70, 192

H

HACH, deep perforating vein of 352
Haemorrhoids 211
Hallux 262, 295, 299
Haustra coli 138–139, 141, 163
HEAD, zones of 67
Heart
- axis 80
- shadow 101
- shape 101
- - aortic knob 101
- - - of pulmonary artery 101
- - - of the left/right atrium 101
- - arch of the ventricle 101
- veins 85
HEISTER's valve 150
Hepar 54, 67, 119–121, 131, 142–146, 149–150, 154–155, 157, 180, 188, 242–244, 246–250, 255
- AP-radiograph 171
- arteries, variations 170
- form variations 144
- laparoscopy 149
- microscopic structure 145
- porta hepatis 142
- projection 144
- segments 147
- veins, projection 146
- - ultrasound image 148
Hiatus
- adductorius 315–316, 323, 358, 362
- aorticus 62–63, 104, 179, T21
- oesophageus 62–63, 104–105, 107, 109, 179, 217, 221, T21
- sacralis 10–11
- saphenus 306, 354–355
- urogenitalis et ani 230
Hilum
- (Glandula suprarenalis) 182
- renale 182
- splenicum 156, 161, 249
Hip
- arteries 359
- muscles 310–322, T42–T43
Horseshoe kidney 185
Humerus 3, 124
HUNTER's vein 352
Hydatid, stalked 201
Hymen 235
Hysterosalpingography 203

I

Ileum 137–140, 157–158, 162–163, 166, 175, 209, 215, 242–244, 251, 254–255
Impotentia
- coeundi 222
- generandi 222
Impressio
- cardiaca 95
- colica 142
- duodenalis 142
- gastrica 142
- oesophagea 142
- renalis 142
- suprarenalis 142
Incisura
- acetabuli 266–267
- angularis 128, 130, 132, 135, 160
- apicis cordis 84–85
- cardiaca pulmonis sinistri 94–95, 103, 110
- cardialis 128, 130
- clavicularis 50
- costalis 50
- fibularis 282
- ischiadica major 266–267
- - minor 266–267, 280
- jugularis (Sternum) 46, 50
- ligamenti teretis 142
- pancreatis 152
- vertebralis inferior 8, 12, 15
- - superior 12, 15
Infundibulum
- ductus paramesonephrici 192
- tubae uterinae 200–201, 204, 209, 228, 253
Inguinal canal 61
- walls 60
Inguinal hernia 61
- direct, medial 61
- indirect, lateral 61
Injection, ventrogluteal 364
Insula pancreatica 153
Interacetabular line 265
Interarticular portion 8, 15
Intermetatarsal joints s. Articulatio(-nes) intermetatarsales
Intersectio tendinea 46, 56–58
Interspinal line 265
Intestinal development 157
Intestinum
- crassum 67, 163–165
- tenue 67, 136, 159, 163–164, 252–253
- - layers of the wall 136
Introitus vaginae 223
Isthmus 8, 15
- glandulae thyroideae 104
- tubae uterinae 201, 209, 229, 253
- uteri 201–202, 253

J

Jejunum 132, 134–135, 137, 152, 154, 157, 162–163, 165, 175, 179, 242, 244, 247, 249–251, 255
Joint
- of CHOPART 296, T39
- of LISFRANC 296, T39
Junctio anorectalis 210
Juncturae cinguli pelvici 268–271, T39

K

KÖHLER's teardrop 280
KOHLRAUSCH's fold 210, 252
KRISTELLER, mucous plug of 207

L

Labium
- anterius 202, 205
- externum 264, 266–267, 322
- internum 264, 266–267
- laterale 275, 277
- majus pudendi 207, 231, 235, 253, 261
- mediale 275–277
- minus pudendi 207, 223, 231, 233, 235, 237, 241, 253, 261
- posterius 202, 205
Labrum acetabuli 279
Lacuna(-ae)
- musculorum 269, 355
- urethrales 198
- vasorum 62, 68, 269, 355
Lamina(-ae)
- anterior (Vagina musculi recti abdominis) 52, 56–58, 70, 72, 355
- arcus vertebrae 7–9, 21–23, 25, 29, 35, 41, 44
- cartilaginis cricoideae 12, 91, 119
- cribrosa 68
- dextra (Cartilago thyroidea) 90–91, 104
- hepatocyticae 145
- muscularis mucosae 105, 130, 136, 139
- parietalis (Pericardium serosum) 74–75, 85, 89, 119–120
- - (Tunica vaginalis testis) 70, 193–194, 239
- posterior (Vagina musculi recti abdominis) 58, 69–70
- profunda (Fascia thoracolumbalis) 29, 32
- propria mucosae 105, 130, 136, 139
- superficialis (Fascia cervicalis) 110, 119
- - (Fascia thoracolumbalis) 29
- visceralis (Pericardium serosum) 74–75, 78–80, 119–120
- - (Tunica vaginalis testis) 194, 239, 242
v. LANZ's point 140
Laparoscopy 149
Larynx 35, 90–91, 109
LAUENSTEIN projection 280
Leg s. Crus
Lien s. Splen
Ligament of GIMBERNAT 355

Ligamentum(-a)
- alare 18
- anococcygeum s. Corpus anococcygeum
- anularia 90–91
- apicis dentis 18–19
- arcuatum laterale 62–63, 68
- - mediale 62–63
- - medianum 63
- arteriosum 74–75, 84, 88, 99, 104–105, 113–114
- atlantooccipitale laterale 17
- bifurcatum 300, 303
- calcaneocuboideum 300, 303
- - plantare 302
- calcaneofibulare 300–304
- calcaneonaviculare 300, 303
- - plantare 299–300, 302–303
- capitis costae intraarticulare 22
- - - radiatum 20–21
- - femoris 236, 256, 260, 271, 279
- - fibulae anterius 286, 294
- - - posterius 287, 290
- cardinale 201, 205
- collaterale(-ia)
- - (Articulationes interphalangeae pedis) 302
- - fibulare 285–289, 293, 310–311
- - laterale 288
- - mediale (Articulatio genus) 287–288
- - - (Articulatio talocruralis) 300–304, 382
- - tibiale 284–290, 293
- coronarium 142–143, 145, 160, 167
- costoclaviculare 51
- costotransversarium 20–21, 32
- - laterale 20–22
- - superius 21–22, 31
- costoxiphoidea 56, 58
- cricothyroideum medianum 90
- cruciatum anterius 286–287, 289–290, 292–293
- - posterius 286–290, 293
- cruciforme atlantis 18–19
- cuboideonaviculare dorsale 300
- - plantare 302
- cuneocuboideum dorsale 303
- cuneonavicularia dorsalia 300, 303
- - plantaria 302
- deltoideum s. Ligamentum collaterale mediale (Articulatio talocruralis)
- denticulatum 40, 43, 45
- epididymidis inferius 193
- - superius 193
- falciforme 59, 71–72, 142–143, 147, 150, 158–160, 166, 243, 248–249
- flavum 20–23, 25, 29, 32, 35, 41, 43–45, 254
- fundiforme penis 58, 60, 238, 242–244, 252
- gastrocolicum 157–158, 160–161, 166, 173, 243, 247
- gastrophrenicum 167
- gastrosplenicum 156, 160–161, 166–167, 245, 247
- hepatoduodenale 131, 150, 160–161, 166–167
- hepatogastricum 131, 157, 160–161, 166, 247, 249
- iliofemorale 268–269, 278–279, 315, 380

Ligamentum(-a)
- iliolumbale 31, 41, 217, 268–270, 272
- infundibulopelvicum 209
- inguinale 57, 60, 62, 69, 196–197, 234–235, 262, 268–270, 306, 313, 317, 354–356, 359
- interclaviculare 51
- intercuneiformia dorsalia 303
- interfoveolare 71
- interspinalia 25, 31, 35, 44, 119
- intertransversarium 21–22, 31–32, 41
- ischiofemorale 278–279
- lacunare 62, 317, 355
- latum uteri 201, 209
- longitudinale anterius 17, 20–22, 24–25, 29, 42, 45, 55, 72, 116, 124, 268
- - posterius 20, 23–25, 42–43, 45
- lumbocostale 63
- meniscofemorale posterius 287–288
- metatarsale(-ia) dorsalia 300
- - plantaria 302
- - transversum profundum 300, 302
- - - superficiale 338
- nuchae 25, 28, 34
- ovarii proprium 200–201, 204, 209
- patellae 285, 287–290, 293, 306, 310, 312–315, 322–323, 327, 332, 356, 368
- pectineum 313, 355
- phrenicocolicum 160, 166
- phrenicosplenicum 247
- plantare(-ia) 302
- - longum 299–300, 302, 304, 342, 379, 383
- popliteum arcuatum 286, 288, 290, 329
- - obliquum 286–288, 293, 329–330
- pubicum inferius 230, 234, 268, 270–272, T39
- - superius 234, 256–257, 268–269, 272, 314, 355, T39
- pubofemorale 278–279
- puboprostaticum 226, 252
- pulmonale 95, 112
- rectouterinum 205
- reflexum 56, 58, 60
- rotundum 167, 201, 204, 229
- sacrococcygeum anterius 230–231
- - posterius profundum 270
- - - superficiale 252, 270
- sacroiliaca anteriora 230, 268–269, 271–272, 317, T39
- - interossea 230, 254, 270–272, T39
- - posteriora 270–271, T39
- sacrospinale 224–225, 229, 240, 265, 269–271, 278, 310, 314, 360, T39
- sacrotuberale 31, 224, 230, 232–233, 240, 265, 269–271, 278, 317, 319–321, 360, 364, T39
- sacrouterinum 201, 253
- splenorenale 156, 245
- sternoclaviculare anterius 51
- sternocostale intraarticulare 51
- - radiatum 51
- supraspinale 22, 25, 31, 33, 270

Ligamentum(-a)
- suspensorium clitoridis 235
- - ovarii 167, 201, 204, 209, 253
- - penis 58, 238–239
- talocalcaneum interosseum 300, 303–304, 383
- - laterale 300
- - mediale 301
- - posterius 301
- talofibulare anterius 300, 303
- - posterius 301, 304
- tarsi plantaria 302
- tarsometatarsalia dorsalia 300, 303, 334
- - plantaria 300, 302
- teres hepatis 59, 71–72, 142, 144, 147, 154, 158, 160–161, 166, 170, 172, 249–250, 255
- - uteri 167, 200–201, 204, 209, 228–229, 253
- tibiofibulare anterius 294, 300–301, 334
- - posterius 300–301
- transversum 23
- - acetabuli 230, 279
- - (Articulatio zygapophysialis) 23
- - atlantis 18–19
- - cervicis s. Ligamentum cardinale
- - genus 286, 289
- - perinei 230, 234
- triangulare dextrum 142
- - sinistrum 142, 147, 167, 244–245
- umbilicale medianum 61, 72, 158, 190–191, 196, 207, 229, 236, 252–253
- venae cavae 142–143, 147
- venosum 142, 147
- vocale 119

Limb, lower 263
- arteries 351
- compartments 366
- cross-section 381–382
- cutaneous innervation, segmental 346
- cutaneous nerves 349
- dermatomes 346
- innervation, segmental, indicator muscles T41
- lymphatics 352
- muscles 306–309, T42–T54
- - origins and insertions 310–311, 324–325
- nerves 345
- skeleton 263
- surface anatomy 262
- veins 352

Limbus
- acetabuli 256, 266–267
- fossae ovalis 78

Linea(-ae)
- alba 46, 56, 58, 60, 72–73, 207, 242–243, 248–249, 251–254
- anocutanea 210
- arcuata 71, 271, 273
- - (Os ilium) 264, 266
- - (Vagina musculi recti abdominis) 58, 70, 158
- aspera 275–277, 323
- axillaris posterior 1
- epiphysialis 281, 291, 293, 304–305, 383
- glutea anterior 267
- - inferior 267
- - posterior 267

Linea(-ae)
- intercondylaris 275
- intermedia 264, 266–267
- intertrochanterica 275
- mediana posterior 1
- musculi solei 282–283
- paravertebralis 1
- pectinea 275–276
- scapularis 1
- semilunaris 58
- supracondylaris lateralis 275
- - medialis 275
- terminalis 167, 230, 236, 264–265, 271, 280
- transversa 10

Lingua 12
Lingula pulmonis sinistri 94–95, 110
LISFRANC's joint 296, T39
Liver puncture 144
Lobar bronchi 92
Lobulus(-i)
- glandulae mammariae 64
- testis 193–194, 252
- thymi 111

Lobus(-i)
- caudatus 142–143, 154, 161, 166, 172, 244, 249
- (Glandula thyroidea) 104–105, 107, 110
- glandulae mammariae 52
- hepatis dexter 131, 140, 142–144, 149–150, 158–161, 163, 166, 172, 188, 244–246, 248–250, 255
- - sinister 120, 131, 142–144, 149–150, 154, 158–161, 166, 170, 172, 242–243, 245, 247–250, 255
- inferior 94–96, 98–99, 103, 110, 118, 121, 124–127, 148, 159, 243, 245–247
- medius (Prostata) 195
- - pulmonis dextri 94–96, 98, 103, 110, 118, 126–127, 159, 245
- prostatae dexter 195
- - sinister 195
- pyramidalis 110
- quadratus 142, 160–161, 172, 249
- superior 94–96, 98–99, 103, 110–111, 118, 124–127, 159, 245
- (Thymus) 111

Lower abdomen, vessels 175, 177
Lower limb s. Limb, lower
Lumbar part of the vertebral column 43
- AP-radiograph 16
- CT 35
- lateral radiograph 15
- ligaments 23
- MRI 24
- vertebral joints 23

Lumbar puncture 44
Lunula valvulae semilunaris 79
Lymphatics, AP-radiograph 219
Lymphography 219

M

Malleolus
- lateralis 262, 283, 294, 300–301, 304–305, 327–328, 332, 334, 336–338, 367, 370–371, 373

Malleolus
- medialis 262, 282, 294, 301, 304–305, 307, 327–330, 334, 336, 338, 371, 373
- tertius 305
Mamma 64–65, 127
- lymphatic drainage 65
- radiography 65
Mammography 65
Manubrium sterni 3, 47, 50–51, 55, 69, 108, 119, 124
Margo
- acetabuli s. Limbus acetabuli
- anterior (Fibula) 283
- - (Pulmo) 94–95, 110–111
- - (Testis) 193
- - (Tibia) 262, 282–283, 306
- falciformis 306, 355
- inferior (Hepar) 142, 145, 163
- - (Pulmo) 94–95, 110
- - (Splen) 156
- interosseus (Fibula) 283, 330–331
- - (Tibia) 282–283
- lateralis (Ren) 182, 216
- liber (Ovarium) 201
- medialis (Glandula suprarenalis) 182, 189
- - (Ren) 182, 216
- - (Scapula) 2
- - (Tibia) 282–283
- mesovaricus 201, 209
- posterior (Fibula) 283
- - (Testis) 193
- superior (Glandula suprarenalis) 182, 189
- - (Splen) 156, 161, 166–167
Massa lateralis atlantis 6, 17, 19
MAY's vein 352, 365
MCBURNEY's point 140
MECKEL's diverticulum 137
Mediastinum 112–113, 116, 119
- anterius 119
- inferius 119
- medium 119
- organisation 119
- posterius 119, 121
- superius 119
- testis 193–194, 252
Medulla
- (Glandula suprarenalis) 189
- renalis 183–184, 188, 246–247, 250
- spinalis 40–41, 119, 123, 222–223, 248–249
Membrana
- atlantooccipitalis anterior 17, 19
- - posterior 17, 19, 34
- intercostalis interna 31, 55
- interossea cruris 283, 286, 294, 301, 366, 381
- obturatoria 236, 260–261, 268, 271, 278–279, 355
- perinei 252
- tectoria 18–19
Membrum
- inferius 206, 263
- - surface anatomy 262
Meniscus(-i)
- (Articulatio genus) 289
- lateralis 286–290, 292–293
- medialis 286–290
Mesenterium 136–137, 164–166, 252
- diverticuli 137
Mesoappendix 163–167
Mesocolon
- sigmoideum 164, 166–167, 179, 217–218, 226
- transversum 139, 157, 160–161, 163, 165–167, 173, 175, 177
Mesosalpinx 201, 209
Metaphysis 304
Metatarsus 263, 298
Mons pubis 235
Motion segment
- cervical 25
- lumbar 25
Mucous plug of KRISTELLER 207
MÜLLERian duct 192, 200
Muscles
- origins and insertions of
- of the abdominal wall 52, 54, 56–58, 60–62, 66, 72–73
- - Crus 310–311, 324–325
- - CT 73
- - Femur 310–311
- - Pelvis 310–311
- - Pes 339
Muscles of the back 26–34
- autochthonous 28, 30–33, T20
- - lateral tract 29–30, 34, T20
- - medial tract 29–30, T20
- - suboccipital muscles 32–34
- deep 28–34
- superficial 26–27
Musculus(-i) 308, 312, 319, 1040–1041, 1045, 1052
- abdominis 52, 56–58, 62, 66, 72–73, 188
- - CT 73
- abductor digiti minimi 304, 333–334, 336–337, 339–342, 377–378, 384, T40, T54
- - hallucis 299, 304, 333, 335, 338–342, 377–378, 384, T40–T41, T52
- adductor(-es) 243, 259
- - brevis 244, 261, 310–312, 315–316, 358, 380, T40–T41, T45
- - hallucis 304, 339–342, 378, 383–384, T40–T41, T52
- - longus 244, 261, 308, 310–317, 354, 356–358, 380–381, T40–T41, T45
- - magnus 260, 286–288, 290, 310–312, 315–317, 319–321, 323, 358, 363–364, 380–381, T40–T41, T45
- - minimus 260, 316, 319–321, T40
- articularis genus 285, 310, T44
- arytenoideus transversus 119
- biceps brachii 52, 54, 122
- - femoris 258, 284, 286, 288, 309–311, 317–324, 328–332, 361–363, 369–370, 380–381, T40–T41, T46
- brachialis 52
- bronchooesophageus 99
- bulbospongiosus 222–223, 226, 232–233, 235, 240–242, 260–261, T22
- coccygeus s. Musculus ischiococcygeus
- colli 34
- constrictor pharyngis inferior 104, 120
- coracobrachialis 54, 122
- cremaster 56–58, 60–61, 70, 193–194, 197, 226, 239, 242–243, 252
Musculus(-i)
- cricothyroideus 104, 110, 120
- dartos 70, 226, 232, 239, 252
- deltoideus 1, 37, 46, 52, 54, 58, 122–123
- digastricus 31, 33–34
- dorsi 26–33, 35, 41
- - CT 29
- - lateral tract 34
- epicranius 39
- erector spinae 1, 27–29, 32, 73, 122, 125, 188, 243, 246–251, 253–254, 319
- extensor digitorum brevis 327, 332–337, 368, 374, 379, 382, 384, T40–T41, T51
- - - longus 306, 308, 310, 324, 326–327, 332–337, 367–368, 374, 379, 381–384, T40–T41, T47
- - hallucis brevis 306, 327, 332–337, 374, 384, T40–T41, T51
- - - longus 304, 306, 308, 324, 326–327, 332–337, 366–368, 374, 379, 381–382, 384, T40–T41, T47
- extremitatis inferioris 308–309
- fibularis brevis 300, 302–304, 324–325, 327, 331–337, 366, 368, 370, 372, 381–382, T40–T41, T48
- - longus 302, 304, 308, 310, 324–327, 329–330, 332, 335, 339, 341–342, 366–368, 370, 372, 381–383, T40–T41, T48
- - tertius 327, 332–337, 366, 368, 382, T47
- fibularis [peroneus] brevis 326
- flexor(-es) accessorius s. Musculus quadratus plantae
- - digiti minimi brevis (Pes) 339–342, 384, T40–T41, T54
- - digitorum 379
- - - brevis 304, 335, 339–342, 377–378, 383–384, T40–T41, T53
- - - longus 304, 311, 325–326, 329–331, 339, 341–342, 366, 371–372, 377–378, 381–384, T40–T41, T50
- - hallucis brevis 339–342, 377–378, 384, T40–T41, T52
- - - longus 304, 325–326, 329–331, 339–342, 366, 371–372, 377–379, 381–384, T40–T41, T50
- gastrocnemius 262, 284, 286–288, 293, 307–309, 311, 320–323, 326–332, 361–362, 366, 369–370, 381, T41, T49
- gemellus inferior 311, 318–321, 363–364, 380, T43
- - superior 311, 318–321, 363–364, 380, T43
- gluteus maximus 1, 37, 52, 215, 226, 232–233, 240–241, 243, 254, 256–257, 259, 262, 279, 309, 311, 317–322, 361–364, 380, T40–T41, T43
- - medius 72, 254, 256, 259, 279, 309, 311, 314–316, 318–321, 361, 363–364, 380, T40–T41, T43
- - - injection 364
- - minimus 256, 279, 310–311, 318, 320–321, 364, 380, T40–T41, T43
Musculus(-i)
- gracilis 232–233, 261, 284, 308, 310–317, 319–321, 323–324, 328, 356–358, 361–362, 369, 381, T40–T41, T45
- iliacus 62–63, 71–72, 217, 220, 231, 244, 254, 257, 259, 308, 310, 312–314, 317, 348, 355–358, T42
- iliococcygeus 230, 243, T22
- iliocostalis cervicis 28, 30, 33–34, T20
- - lumborum 28, 30, 72, T20
- - thoracis 28, 30, 37, 249, T20
- iliopsoas 71, 73, 256–257, 260–261, 279, 308, 310–316, 321, 355, 380, T40–T41
- infraspinatus 26–28, 37, 52, 123–125
- intercostales 55, 72, 217, 248
- - externi 31, 54–55, 57–58, 60, 106, 115, 124, 126–127, 144, 249, T13
- - interni 31, 54–58, 60, 106, 115–116, 120, 122, 144, 249, T13
- interossei, compartments 379
- - dorsales (Pes) 304, 333–334, 336–337, 339, 341–343, 383–384, T53
- - plantares 333, 339–343, 384, T53
- interspinales cervicis 30–31, 33, 40, T20
- - lumborum 30–31, T20
- - thoracis 30, 119, T20
- intertransversarii anteriores cervicis 33, T20
- - laterales lumborum 31–32, 41, T20
- - mediales lumborum 31–32, 41, T20
- - posteriores cervicis 31, 34, T20
- - - lumborum 30
- - thoracis 34, T20
- ischiocavernosus 222–223, 226, 232–233, 235, 237, 240–242, 260–261, T22
- ischiococcygeus 225, 227, 230–231, 256, 317, T22, T40
- ischiocrurales 260
- latissimus dorsi 1, 26–29, 37–38, 52, 54, 57–58, 63, 72, 126–127, 248–251, 319, T18
- levator(-es) ani 210, 212–214, 226–233, 236–237, 240–243, 256–258, 260–261, 311, T22, T40
- - costarum T20
- - - breves 31–32
- - - longi 31–32, T20
- - scapulae 27–28, 34, 37–38, T17
- longissimus capitis 28, 30–31, 33–34, 37–39, T20
- - cervicis 28, 30, 33–34, T20
- - thoracis 28, 30, 37, 72, 249, T20
- longus colli 55, T12–T13
- lumbricales (Pes) 340–342, T40, T53
- multifidus(-i) cervicis 30, 33–34, 39, 1020
- - lumborum 30–32, 249, T20
- - thoracis 30–32, 37, T20
- obliquus capitis inferior 31, 33, 39–40, T20
- - - superior 31, 33, 39–40, T20

Musculus(-i) obliquus
- - externus abdominis 26–29, 31, 37, 46, 52, 54, 56–58, 60–61, 63, 66, 68, 70–73, 110, 158, 217, 231, 239, 248, 251, 254, 311, 322, 355, 364, T15, T40
- - internus abdominis 27–28, 31, 56–58, 60–61, 66, 68, 70–73, 158, 217, 231, 239, 251, 254, 310, T15, T40
- obturatorius externus 236, 256–257, 259–261, 279, 310–311, 316, 321, 364, 380, T40–T41, T45
- - internus 212–213, 225, 230–231, 236–237, 256–261, 279, 310–311, 317–321, 363–364, 380, T40–T41, T43
- occipitofrontalis 39
- omohyoideus 110, 123
- opponens digiti minimi (Pes) 334, 342, 384, T40, T54
- papillaris anterior 78–82
- - posterior 78–81
- - septalis 78
- pectinati 78
- pectineus 244, 256–257, 308, 310–316, 355–357, 380, T40–T41, T45
- pectoralis major 46, 52, 54, 56–58, 60, 64, 110, 119, 123–127, T26
- - minor 54, 60, 120, 124, T26
- perinei 230
- peroneus brevis, longus bzw. tertius s. Musculus fibularis brevis, longus bzw. tertius
- pharyngis 109
- piriformis 224, 226–227, 231, 243, 310–311, 314–321, 360, 363–364, 380, T40–T41, T43
- plantaris 286, 311, 323, 328–331, 366, 369–372, 382, T40, T49
- popliteus 286–290, 293, 311, 325, 330–331, 369–370, 372, T40–T41, T50
- psoas major 29, 35, 41, 62–63, 68, 72, 152, 185–186, 188, 217, 221, 236, 245–247, 251, 254, 257, 259, 308, 310, 312–314, 317, 348, 354–355, T42
- - minor 62–63, 72, 179, 217, 313, 317, 355, T42
- pubococcygeus 230, 242–243, T22
- puborectalis 230, 256, 258
- pyramidalis 56–58, 66, 70, 72, 242, T14
- quadratus femoris 260, 310, 318–321, 363–364, 380, T40, T43
- - lumborum 29, 31, 62–63, 66, 68, 72–73, 121, 152, 217, 220–221, 246–247, 251, 310, T16
- - plantae 304, 339, 341–342, 377–378, 382–383, T40, T53
- quadriceps femoris 262, 285, 287–288, 290, 306, 308, 311–312, 314, 324, 332, T40–T41, T44
- rectococcygeus 230–231
- rectouterinus 201
- rectus abdominis 46, 56–59, 61–63, 66, 69–73, 110, 119, 155, 158, 217, 226, 242–243, 246–254, 256, T14, T40

Musculus(-i) rectus
- - capitis anterior T12
- - - lateralis 34, T20
- - - posterior major 31, 33, 39–40, T20
- - - - minor 31, 33, 40, T20
- - femoris 244, 256–257, 269, 278–279, 308, 310–317, 322, 356–358, 380–381, T44
- rhomboideus major 26–28, 37–39, T17
- - minor 27, 38–39, T17
- rotatores breves 30–31, T20
- - longi 30–31, T20
- sacrococcygeus 230–231, 243
- sartorius 52, 244, 256–257, 262, 284, 308, 310–317, 322–324, 356–358, 362, 380–381, T40–T41, T44
- scalenus anterior 55, 110, 112, 117–118, 123, T11
- - medius 34, 40, 55, 120, 122, T11
- - posterior 28, 31, 34, 55, T11
- semimembranosus 258, 284, 286–288, 309–311, 317–318, 320–323, 325, 328–330, 361–363, 369–370, 380–381, T41, T46
- semispinalis capitis 28, 30–31, 33–34, 38–40, T20
- - cervicis 28, 30–31, 33, 39, T20
- - thoracis 31, 33–34, T20
- semitendinosus 258, 284, 309–311, 316–321, 323–324, 328, 361–363, 369–370, 380–381, T41, T46
- serratus anterior 27–28, 46, 52, 54, 56–58, 63–64, 70, 120, 122–127, T17
- - posterior inferior 27–28, 37, 249, T19
- - - superior 27–28, 37, T19
- soleus 290, 308–309, 311, 325–332, 369–372, 381, T41, T49
- sphincter ani externus 207, 210, 212–214, 226, 230–233, 235, 240–242, 252, 258, T22, T40
- - - internus 207, 210, 231, 242, 252, 258
- - pyloricus 135, 152
- - urethrae 207, 226, 231, 234, T22
- - vesicae 222
- spinalis 30
- - capitis 30–31, T20
- - cervicis 30, T20
- - thoracis 28, 30, T20
- splenius capitis 26–28, 30–31, 33–34, 38–39, 122, T20
- - cervicis 27–28, 30, 33–34, T20
- sternalis 54, 60, T13
- sternocleidomastoideus 26–28, 35, 38, 54, 58, 110, 119, 123
- sternohyoideus 110, 123
- sternothyroideus 55, 110, 119, 124
- subclavius 54, 60, 110, 112–113, 120, 122–123, T26
- subcostales 60, T13
- suboccipitales 31–34, 40
- subscapularis 60, 122, 124, T26
- supraspinatus 122–123
- suspensorius duodeni 135, 152
- tensor fasciae latae 52, 73, 244, 256–257, 262, 279, 306–308, 311–314, 320, 322, 356, 380, T40–T41, T44

Musculus(-i)
- teres major 1, 26–28, 37, 52, 122, 124–125
- - minor 37, 52
- thoracis 31, 52, 54, 60
- thyrohyoideus 110
- tibialis anterior 290, 300, 302, 306, 308, 310, 324, 326–327, 332, 334, 336–337, 339, 366–368, 374, 381–383, T40–T41, T47
- - posterior 300, 302, 304, 311, 325–326, 329–331, 339, 342, 366, 370–372, 381–382, T40–T41, T50
- trachealis 91
- transversospinalis 29
- transversus abdominis 31, 58–63, 66, 68, 70–73, 158, 217, 221, 231, 249, 251, 254, 310, T15, T40
- - perinei profundus 207, 226, 230, 232–236, 242, 252, 260, 311, T22, 1040 Tab
- - - superficialis 230, 232–235, 240–241, 260, T22, 1040 Tab
- - thoracis 55, 69, 110, 127, T13–T14
- trapezius 1, 26–28, 33–34, 38, 52, 122–126, T17
- triceps brachii 37, 52
- - surae 304, 326, 333, 366, 383, T40, T49
- vastus intermedius 310–312, 315–316, 381, T44
- - lateralis 257, 259, 279, 290, 293, 308, 310–316, 320–323, 332, 356, 358, 361, 381, T44
- - medialis 290, 308, 310–316, 323, 356–358, 381, T44

Myocardium 76, 78–80, 119

N

Neck, muscles 34
Neck-shaft angle of the femur s. Angle between neck and shaft of the femur
Nephros s. Ren
Nervous system, autonomous 117, 120–121
Nervus(-i)
- accessorius [XI] 38–39, T17
- anales 214
- - inferiores 225, 240–241, 344, T40
- anococcygeus 225, 240–241, 344, T40
- auricularis magnus 36–38
- axillaris 36–37, 122
- cardiacus cervicalis inferior 83
- - - medius 83, 120
- - - superior 83
- cavernosi clitoridis 223
- cervicalis 19, 40, 120, T19–T20
- clunium inferiores 36, 240–241, 349, 360–362, T40
- - medii 36, 349, 360–362
- - superiores 36–37, 349, 360, 362
- coccygeus 225, 344, T40
- cutaneus brachii lateralis inferior 36–37
- - - - superior 36
- - - posterior 36–37

Nervus(-i) cutaneus
- - dorsalis intermedius 345, 349, 365, 373, T40
- - - lateralis 345, 349–350, 365, 373, T40
- - - medialis 345, 349–350, 365
- - femoris lateralis 36, 68, 71, 73, 179, 217, 220–221, 317, 344–345, 348–349, 353, 355–357, 360, T40
- - - posterior 36, 240–241, 257, 344–345, 349, 360–364, 380, T40
- - surae lateralis 345, 349–350, 361–363, 365, 369, T40
- - - medialis 345, 361–363, 365–366, 369, 373, T40
- digitales dorsales pedis 345, 349, 368, 373–374, T40
- - plantares communes 376–378, T40
- - - proprii 376, 378, T40
- dorsalis clitoridis 223, 234, 241, T40
- - penis 222, 234, 238–240, T40
- - scapulae 38–39, T17
- erigentes s. Nervus(-i) splanchnici pelvici
- femoralis 66, 68, 73, 179, 217–218, 220–221, 236, 244, 256–257, 317, 344–345, 348–349, 353, 355–358, 380–381, T40, T44–T45
- fibularis communis 284, 344–345, 349–350, 361–363, 365, 368–371, T40, T51
- - profundus 345, 349–350, 366, 368, 373–374, 379, 381–382, T40, T47, T51
- - superficialis 345, 349–350, 365–366, 368, 373, 379, 381–382, T40, T48
- genitofemoralis 60, 66, 68, 179, 197, 217, 220–221, 225, 229, 236, 238, 256, 317, 344–345, 348–349, 353, 355, T15, T40
- gluteus inferior 257, 344–345, 349, 360, 363–364, 380, T40, T43
- - superior 344–345, 349, 360, 364, 380, T40, T43–T44
- hypogastricus 214, 222–223
- iliohypogastricus 36–37, 66, 68, 73, 115, 117, 179, 217, 220–221, 344–345, 348–349, 353, 364, T15, T40
- ilioinguinalis 60, 66, 68, 73, 179, 217–218, 220–221, 238–239, 344–345, 348–349, 353, T15, T40
- intercostales 36, 38, 41, 54, 66, 70, 73, 112–113, 115–117, 120, 124, 127, 220, 247–248, T13–T17
- intercostobrachialis 37
- interosseus cruris T40
- ischiadicus 256–257, 284, 344–345, 347, 349–350, 360, 362–364, 380–381, T40, T43, T46–T50
- - projection 360
- labiales 223
- - anteriores T40
- - posteriores 223, 241, T40
- laryngeus recurrens 83, 88–89, 111–113, 115, 117–118, 120
- lumbales 66, 226, 349

Nervus(-i)
- obturatorius 68, 71, 220–221, 225, 229, 231, 236–237, 256–257, 261, 317, 344–345, 348–349, 353, 355, 357–358, 360, 380, T40, T45
- occipitalis major 37–40
- - minor 36–38
- - tertius 37, 40
- pectoralis lateralis 60, T26
- - medialis 60, T26
- perineales 234, 240–241, 260–261, T40
- phrenicus 63, 89, 110–113, 115, 118, 120–121, 124–127, T21
- - course 118
- plantaris lateralis 304, 345, 350, 376–378, 382–383, T40, T52–T54
- - medialis 304, 345, 350, 376–378, 382, T40, T52–T53
- pudendus 214, 222–223, 225, 234, 236, 240–241, 257, 260–261, 344–345, 349, 360, T22, T40
- radialis 36–37
- rectales inferiores s. Nervus(-i) anales inferiores
- sacrales 223, 225–226, 243, 349, T22
- saphenus 345, 348–349, 353, 357–358, 365–366, 373, 381–382, T40
- scrotales anteriores T40
- - posteriores 222, 240, T40
- spinales 36, 41, 43, 45, 251
- spinalis 24, 29
- spinalis(-es) 24, 35, 40–41
- splanchnici lumbales 214
- - pelvici s. Radix parasympathica
- splanchnicus major 63, 112–113, 115–117, 120–121, 220–221, 245, 249
- - minor 63, 115, 117, 121, 245, 249
- subcostalis 68, 73, 115–117, 217, 220–221, 344
- suboccipitalis 37, 39–40, T20
- subscapulares 37, T26
- supraclaviculares 66
- - laterales 36–37
- suprascapularis 123
- suralis 345, 349–350, 362, 365, 382, T40
- thoracici 37, 60, 66, 70, 112–113, 115–117, 120–121, 220, 249, T15, T19–T20
- thoracicus longus 122, 124–126, T17
- thoracodorsalis 122, T18
- tibialis 284, 344–345, 349–350, 361–363, 365–366, 369–373, 376, 381–382, T40, T49–T50, T52–T53
- vaginalis 223
- vagus [X] 83, 88–89, 111–113, 115, 117–118, 120–121, 123–127, 220

Node of ROSENMÜLLER 355

Nodulus(-i)
- lymphoideus(-i) aggregati 137, 139
- - solitarii 130, 136–137, 139, 150, 210
- valvulae semilunaris 79

Nodus(-i)
- atrioventricularis 82–83
- lymphoideus(-i) aortici laterales 229
- - axillaris(-es) 65, 122, 124
- - - apicalis 124

Nodus(-i) lymphoideus(-i)
- - brachiales 65
- - bronchopulmonales 125–127
- - cervicales anteriores 65
- - - - profundi 65
- - - laterales 120
- - - - profundi inferiores 120
- - deltopectorales 65
- - gastrici 131, 218
- - gastroomentales 131, 249
- - hepatici 131
- - ileocolici 243
- - iliaci communes 218, 229
- - - externi 218, 229, 236, 354
- - - - laterales 229, 354
- - - - mediales 229, 354
- - - interni 218, 229, 354
- - infraclaviculares 65
- - inguinales 352
- - - profundi 218, 354
- - - superficiales 218–219, 354
- - - - inferiores 218, 354
- - - - superolaterales 218, 354
- - - - superomediales 218, 256, 354
- - juxtaoesophageales 107, 127
- - lacunaris(-es) 355
- - - medialis 355
- - ligamenti arteriosi 88
- - lumbales dextri 218
- - - intermedii 218
- - mediastinales anteriores 88, 111, 113
- - - posteriores 107, 245
- - obturatorii 229
- - pancreaticus 250
- - paramammarii 65
- - pararectales 122, 229
- - parasternales 65
- - paratracheales 88, 120, 124
- - parietales 218
- - phrenici inferiores 218
- - - superiores 88, 119
- - popliteus profundus 366
- - - superficialis 366
- - preaortici 229, 251
- - precavales 229
- - promontorii 218
- - pylorici 131
- - rectalis superior 229
- - retroaortici 229
- - retrocavales 243
- - splenici 245, 249
- - submammarii 65
- - supraclaviculares 65
- - thoracis 107, 120
- - tibialis anterior 367
- - tracheobronchiales 95, 98–99, 113, 122
- - - inferiores 88, 95, 107, 119, 127
- - - superiores 88, 107, 119
- sinuatrialis 82

Nucleus pulposus 24–25, 29

O

Oesophagoscopy 108, 133

Oesophagus 62–63, 67, 104–109, 112–113, 115–117, 119–121, 123–130, 132–133, 157, 180, 220, 242, 245
- constrictions 108–109
- diverticulum 109
- radiography 108

Olisthesis 9

Omentum
- majus 121, 131, 139–140, 149, 157–166, 172, 242–243, 245–247, 249, 252, 254
- minus 160–161, 166, 247, 249

Orchis s. Testis

Organum(-a)
- genitalia feminina externa 223, 233, 235, 241
- - - interna 204, 209
- - masculina 222, 239
- - - externa 197, 238, 240
- pelvis in the female 237
- - in the male 236
- urogenitalia feminina, development 200
- - masculina, development 192

Orificium ductus pancreatici 150

Os(-sa)
- coccygis 2–4, 11, 207, 215, 230, 232–233, 242, 252, 256–257, 271, 273
- coxae 2, 16, 41, 263, 266–267
- - AP-radiograph 16
- - development 267
- cuboideum 295–296, 298–299, 303, 383
- cuneiforme(-ia) 299
- - intermedium 295–296, 298, 304, 383
- - laterale 295, 298
- - mediale 295–296, 298, 302
- digitorum (Pes) 263, 295–296
- hyoideum 12
- ilium 3, 15–16, 41, 71, 73, 230, 244, 258, 264, 272, 274, 280
- ischii 236, 256, 264, 274
- metatarsi 263, 295–296, 298–299, 303–304, 379, 383
- naviculare 295–296, 298–299, 303–305, 383
- occipitale 6, 12, 17–19
- pedis 295–296, 299
- - structure characteristics 299
- pubis 196, 215, 226, 234, 242, 244, 256–257, 264, 274, 279
- sacrum 2–5, 10–11, 15, 23, 26, 41, 215, 226, 230, 242–243, 252–254, 262, 264, 271–273, 317
- - sex differences 11
- sesamoidea (Pes) 295–296, 302, 339
- tarsi 263, 295–296, 298
- temporale 40

Ostium(-a)
- abdominale tubae uterinae 201, 204, 209
- appendicis vermiformis 139
- atrioventriculare dextrum 78
- - sinistrum 79, 126
- cardiacum 108, 121, 130, 166–167, 244–245, 247
- ileale 139
- pyloricum 135
- sinus coronarii 78, 82, 84
- ureteris 190–191, 195, 198, 205, 252–253, 260
- urethrae externum (Urethra feminina) 200, 231, 233, 235, 237, 241, 253, 261
- - - (Urethra masculina) 198–199, 252
- - internum (Urethra feminina) 205, 253
- - - (Urethra masculina) 190, 195, 198, 252, 260

Ostium(-a)
- uteri 201–202, 205, 207, 253
- uterinum tubae uterinae 201, 203
- vaginae 200, 233, 235, 237
- venae cavae inferioris 82
- - - superioris 82

Ovarium 140, 167, 200–202, 204, 209, 223, 228, 253, 354

P

Palatum osseum 12

Pancreas 121, 131, 134, 152–153, 157, 163–164, 172, 177, 242–243, 245, 249–250
- computed tomography 155
- low magnitude enlargement 153
- projection 134
- ultrasound image 154

Pancreatocyticus exocrinus 153

Panniculus adiposus 58, 72, 210, 232, 239

Papilla(-ae)
- duodeni major 135, 150, 152–153
- - minor 152–153
- ilealis 139
- mammaria 52, 64, 127
- renales 183–184, 186, 245
- of Vater s. Papilla(-ae) duodeni major

Paracystium 260–261

Paradidymis 192

Parametrium s. Ligamentum cardinale

Paraproctium s. Spatium pararectale

Parasympathetic nervous system 214, 222–223

Paries
- anterior (Gaster) 133, 160–161
- membranaceus (Trachea) 91, 99, 107, 119
- posterior (Gaster) 133, 155

Paroophoron 200

Pars
- abdominalis aortae 35, 62–63, 69, 104, 115, 117, 148, 154–155, 157, 168, 172, 174, 177–179, 185, 187–189, 212, 214, 216–218, 221, 225, 227, 229, 242, 245, 250–251, 359
- - - ultrasound image 178
- - autonomica 220
- - (M. pectoralis major) 52, 54, 56, 58, 60, T26
- - (Oesophagus) 62–63, 104, 106, 108–109, 117, 128, 130
- - (Ureter) 163, 216
- anterior (Facies diaphragmatica hepatis) 145
- - (Fornix vaginae) 202, 205
- anularis vaginae fibrosae 340
- ascendens aortae 75, 78–80, 84, 109, 114–115, 117, 119–121, 125–126, 255
- - (Duodenum) 134, 139–140, 152, 165, 167
- - (M. trapezius) 26, T17
- atlantica (A. vertebralis) 34, 40
- basalis (Aa. lobares inferiores dextrae) 98
- - (Aa. lobares inferiores sinistrae) 98–99
- basilaris (Os occipitale) 17–19

Pars
- cardiaca 63, 104, 121, 130–132, 179, 217, 242
- cervicalis (Columna vertebralis) 2
- – (Oesophagus) 104, 106–108, 119
- clavicularis (M. pectoralis major) 54, T26
- costalis diaphragmatis 54, 62–63, 73, 148, 179, 246–250, T21
- – (Pleura parietalis) 54, 110, 112–113, 120–121, 124, 159, 246–250
- cruciformis vaginae fibrosae 340
- descendens aortae 75, 107, 109, 114, 125–127
- – (Duodenum) 128, 130, 132, 134–135, 140, 152–154, 167, 179
- – (Lig. iliofemorale) 278
- – (M. trapezius) 26, T17
- diaphragmatica 54, 63, 89, 110–113, 118, 159, 245, 248–250
- horizontalis (Duodenum) 134–135, 152–153, 163, 165, 167, 179
- infraclavicularis 54, 112–113
- intermedia (Urethra masculina) 195–196, 198, 252
- intramuralis (Urethra masculina) 198
- intrasegmentalis 120, 122
- lateralis (Os occipitale) 17
- – (Os sacrum) 10–11, 41, 254, 264, 273
- libera membri inferioris 263
- lumbalis (Columna vertebralis) 2
- – diaphragmatis 62–63, 68, 104, 106, 115–116, 119, 121, 148, 155, 172, 242–243, 245, 248–250, T21
- mediastinalis 89, 95, 111–113, 125, 159
- membranacea (Septum interventriculare) 79–80
- – (Urethra masculina) s. Pars intermedia (Urethra masculina)
- muscularis (Septum interventriculare) 78–80
- occlusa 229, 236
- patens 225, 227–229
- pelvica (Ureter) 71, 166–167, 221
- posterior (Fornix vaginae) 202, 205, 207, 253
- profunda (Compartimentum cruris posterius) 366
- prostatica 190, 195, 198
- pylorica 117, 130, 132–135, 158, 160, 166, 247, 255
- spongiosa 198–199, 252
- sternalis diaphragmatis 62–63, 110–111, 119, 243, T21
- sternocostalis (M. pectoralis major) 52, 54, 56, 58, 60, T26
- subcutanea (M. sphincter ani externus) 235, T22
- superficialis (Compartimentum cruris posterius) 366
- superior (Duodenum) 128, 130, 134–136, 149, 153, 160–161, 166–167, 179, 244
- – (Facies diaphragmatica hepatis) 142

Pars
- terminalis 137, 163
- thoracica aortae 63, 69, 87, 99, 104–106, 109, 113–115, 117, 121, 124, 152, 157, 178, 248
- – autonomica 117, 120–121
- – (Columna vertebralis) 2
- – (Ductus thoracicus) 113, 115–116
- – (Oesophagus) 63, 104–109, 113, 115–117, 119, 121, 245
- tibiocalcanea 300–302, 304
- tibionavicularis 300
- tibiotalaris anterior 300
- – posterior 300–301
- transversa (Lig. iliofemorale) 278
- – (M. trapezius) 26, T17
- transversaria (A. vertebralis) 40
- uterina (Tuba uterina) 201

Patella 262–263, 284–285, 287, 290–293, 306, 314–315, 322–323, 327, 332, 365
- ridge of the 292

Pecten
- analis 210, 231, 252
- ossis pubis 62, 231, 264, 266–267, 271, 314–315, 317

Pediculus arcus vertebrae 5, 7–9, 12, 14–16, 20–21, 23–25, 29, 35, 273
Pedograms 384
Pelvic aperture 264
Pelvic arteries, digital subtraction angiography (DSA) 359
Pelvic constriction 265
Pelvic diaphragm T22
- of the female 230–231, 233–235
- of the male 232, 234, 236

Pelvic organs
- in the female 228
- in the male 226–227

Pelvic width 265
Pelvis 264, 268
- AP-radiograph 273–274
- CT 254
- of the female 265
- – cross-section 257
- – CT 257
- – median section 253
- – MRI 253, 259
- – oblique section 261
- – pelvic dimensions 265
- joints of pelvic girdle 268–271
- lumbosacral passage 268–269, 271
- major 264
- of the male, cross-section 254, 256
- – CT 256
- – frontal section 258
- – median section 242, 252
- – MRI 252, 259
- – oblique section 260
- – sagittal section 243
- minor 167, 264
- muscles, origins and insertions 310–311
- renalis 171, 182–184, 186, 188, 192, 200, 216, 245–246, 250
- – ampullar type 186
- – AP-radiograph 186
- – dendritic type 186
- sex differences 264, 268–271

Penis 197, 199, 222, 244, 260
Perforating vein, lateral 352
Pericardium 63, 74–75, 85, 89, 98–99, 110, 118, 247
- fibrosum 74, 89, 107, 111–113, 121, 159
- serosum 74–75, 78–80, 84, 89, 119–120

Perineal laceration 233
Perineum 235
- of the female 230, 233, 241
- of the male 232, 240

Periorchium 194
Periosteum 43
Peristalsis 132
Peritoneum 143
- parietale 54, 59, 61–62, 75, 120, 149, 158, 167, 179, 190, 196, 217, 226, 242, 248–251, 253, 258
- urogenitale 202, 210, 260–261
- viscerale 54, 70, 120, 136, 242, 246, 248–249, 251

Pes 263
- anserinus superficialis 313, 323
- cavus 384
- compartments 379
- cross-section 382
- frontal section 384
- joints 300–303, T39
- ligaments 302
- longitudinal section, MRI 383
- muscles 333–343, 384, T51–T54
- – origins and insertions 333, 339
- osteofibrous tubes 384
- sagittal section 383
- tendon sheaths 334–335
- valgus 303
- vault 299

PEYER's patches 137
Phalanx(-ges)
- distalis(-es) (Pes) 263, 295–296, 383
- media(-ae) (Pes) 263, 295–296, 383
- (Pes) 263, 296, 299, 304
- proximalis (Pes) 295–296, 383

Pharynx 133
Pia mater spinalis 41, 43
Placenta 207–208
Planta 262, 375–378
- arteries 375
- – variations 375

Plantar aponeurosis s. Aponeurosis plantaris
Platysma 58
Pleura
- parietalis 54, 63, 89, 95, 110–113, 118, 120–121, 123–125, 159, 245–250
- pulmonalis s. Pleura visceralis
- visceralis 54, 112, 120–124, 246–247

Pleural borders, projection 103
Pleurocentesis, puncture of the pleural cavity 54
Plexus
- aorticus thoracicus 89, 113, 117, 121
- brachialis 54, 60, 111–113, 120, 122–123, 128, T11, T17
- cardiacus 83, 113, 117–118
- caroticus communis 110
- cervicalis 36–37, 40, T11–T12, T17, T21
- coccygeus 225, 344, T40
- coeliacus 214, 220
- deferentialis 197, 222

Plexus
- hypogastricus inferior 222–223
- – superior 214, 222–223
- lumbalis 29, 36, 66, 68, 254, 344, T15, T40, T42, T45
- lumbosacralis 68, 220, 344, T40
- mesentericus inferior 220, 222
- – superior 222–223
- oesophageus 112, 117, 121, 220
- ovaricus 223
- pampiniformis 60, 70, 193–194, 197, 238–239, 243
- prostaticus 222
- pulmonalis 112–113, 120–121
- rectalis inferior 214
- – medius 214
- – superior 214
- renalis 223
- sacralis 68, 214, 220, 223, 225, 344, 349, T22, T40, T43, T46
- testicularis 197, 222
- uterovaginalis 223
- venosus areolaris 66
- – prostaticus 236, 260
- – rectalis 213
- – submucosus 106, 181
- – vaginalis 228
- – vertebralis externus anterior 42
- – – – posterior 42
- – – internus 123
- – – – anterior 41–43, 45, 242
- – – – posterior 41–44, 119
- – vesicalis 227, 236–237, 256

Plica(-ae)
- alares 285
- circulares 130, 132, 135, 137, 150, 152
- duodenalis inferior 165, 167
- gastricae 130, 133
- gastropancreatica 161, 166–167
- ileocaecalis 162–163, 167
- interureterica 190–191, 198, 260
- longitudinalis duodeni 150, 152
- mucosae 150, 190, 198
- palmatae 201, 253
- rectouterina 201, 228–229, 253
- semilunares coli 139, 141, 163
- spiralis 150–151
- synovialis 301
- – infrapatellaris 285
- – mediopatellaris 284
- transversa recti 207, 210–211, 242, 252–253
- tubariae 201
- umbilicalis lateralis 59, 71–72, 158–159, 227, 236
- – medialis 59, 71–72, 158–159, 196, 209, 217, 227, 236, 253
- – mediana 59, 71–72, 158–159, 196, 209, 217, 236, 252–253
- vesicalis transversa 71, 236

Point
- of v. LANZ 140
- of MCBURNEY 140
- of SUDECK 176, 212

Polus urinarius 184
Popliteal perforating vein 352
Porta hepatis 142
Portio
- supravaginalis cervicis 201–202
- vaginalis cervicis 201, 205, 207, 253
- – – nullipara 205

Portocaval anastomoses 181, 213
Positio (Uterus) 203

Preputium
- clitoridis 233, 235
- penis 197–199, 242–243, 252
Processus(-us)
- accessorius 5, 8–9
- articularis inferior 6, 8–9, 12, 15–16, 22–23, 25, 29, 35, 254
- - superior (Os sacrum) 10–11, 41, 230, 254
- - - (Vertebrae) 5–9, 12, 15–16, 20–21, 23, 25, 29, 35, 41, 45, 248, 273
- caudatus 142
- coracoideus 47–48, 122
- costalis 5, 9, 16, 23, 30, 32, 35, 48, 63, 216, 269, 272–273
- falciformis 270
- lateralis tali 295, 297–298, 382
- - tuberis calcanei 295, 297–298
- mamillaris 8–9, 35
- mastoideus 12, 33, 40
- medialis tuberis calcanei 295–298, 302
- papillaris 142, 166, 249
- posterior tali 296–298, 305
- spinosus 1, 5–9, 12–16, 19, 22–26, 28–29, 32–35, 41, 44, 119, 123–124, 127, 188, 249, 251, 273
- styloideus (Os temporale) 33–34
- transversus 6–8, 13–14, 20, 22, 31, 34–35, 40, 125
- - (Atlas) 34, 40
- - (Axis) 34
- uncinatus (Pancreas) 152–153, 173, 179, 243
- - (Vertebra) s. Uncus corporis
- vaginalis peritonei 61
- - - persistens 61
- xiphoideus 47, 50, 55, 63, 70, 119, 127, 132, 207, 242, 248
Promontorium (Os sacrum) 3–4, 10–11, 15, 62, 177, 185, 217, 227, 242, 253, 264, 314
Pronatio (Articulatio talotarsalis) T39
Prostata 71, 190–191, 195–196, 198, 222, 226–227, 236, 242–243, 252, 259–260
Protuberantia occipitalis externa 26, 39
Psoas arcade 63
Pubes 197
Pulmo 54, 92, 94–100, 110
- arteries 100
- dexter 89, 94–98, 110–111, 118, 120–121, 123–127, 148, 159, 243, 245–246
- organisation 98
- sinister 89, 94–99, 110–111, 118, 120–122, 124, 126–127, 159, 245, 247
- veins 100
Pulmonary angiography 100
Pulmonary borders, projection 103
Pulpa splenica 156
Pyelography, intravenous 216
Pylorus 128–131, 135, 160–161
Pyramides renales 183–184, 188, 245

Q

Quadrangular space 37

R

Radix(-ces)
- anterior (N. spinalis) 41, 43, 45, 214
- mesenterii 140, 163, 166–167, 177
- motoria s. Radix anterior (N. spinalis)
- parasympathica 214, 222–223
- posterior (N. spinalis) 41, 43, 45
Ramus(-i)
- acetabularis (A. circumflexa femoris medialis) 359
- - (A. obturatoria) 357, 359
- anterior (A. media genus) 289
- - (A. obturatoria) 357, 359
- - (A. renalis) 182
- - (N. cervicalis) 120
- - (N. lumbalis) 41, 226
- - (N. obturatorius) 68, 348, T39–T40
- - (N. sacralis) 226
- - (N. spinalis) 40–41, 45, T20
- - (Truncus n. spinalis) 42–43, 45
- - (V. portae hepatis) 246
- articulares (A. descendens genus) 351, 356–358
- ascendens (A. circumflexa femoris lateralis) 351, 358–359
- - (A. circumflexa femoris medialis) 363
- atriales (A. coronaria dextra) 84, 86
- - (A. coronaria sinistra) 86
- atrioventriculares 86–87
- bronchiales 99
- - (Aorta) 105–106, 112, 115, 121
- - (N. vagus) 115, 117, 120–121
- calcanei (A. tibialis posterior) 372
- - mediales (N. tibialis) 376, T40
- cardiacus cervicalis inferior 376
- - - - (N. vagus) 83
- - thoracicus inferior (N. vagus) 83, 118
- - - (N. vagus) 112–113, 118
- - - (Truncus sympathicus) 113
- circumflexus (A. coronaria sinistra) 74–75, 84–87
- communicans(-tes) (A. fibularis) 351
- - albus 42
- - fibularis 345, 365, T40
- - griseus 42
- - (Truncus n. spinalis) 41, 43, 45
- - (Truncus sympathicus) 45, 112–113, 116–117, 222
- coni arteriosi 84, 86–87
- cutaneus(-i) anterior(-es) abdominales 66
- - - (N. femoralis) 66, 68, 348–349, 353, T40
- - - (N. iliohypogastricus) 66, 349, 353
- - cruris mediales (N. saphenus) 349, 365, T40
- - lateralis(-es) (N. iliohypogastricus) 36, 349, 360, 364
- - - (N. spinalis) 36, 41
- - - (Nn. thoracici) 344
- - - pectorales (Nn. intercostales) 38, 66

Ramus(-i) cutaneus(-i)
- - medialis(-es) (N. spinalis) 36, 41
- - (N. obturatorius) 349, 353, 357–358, 360, T40
- - (N. plantaris medialis) 377
- - posterior(-es) (Nn. cervicales, Rr. posteriores) 38
- - - (Nn. thoracici, Rr. posteriores) 38
- descendens (A. circumflexa femoris lateralis) 351, 357–359
- - (A. circumflexa femoris medialis) 359
- dexter (A. hepatica propria) 146, 171, 249
- - (V. portae hepatis) 146, 244, 246, 248–249
- dorsalis (A. lumbalis) 41
- - (V. lumbalis) 41
- femoralis 66, 68, 217, 221, 317, 344, 349, 353, 355, T40
- fundi uteri (A. uterina) 204
- gastrici anteriores (Truncus vagalis anterior) 117
- - (V. gastroomentalis dextra) 160
- genitalis 60, 68, 197, 217, 221, 225, 238, 256, 344, 349, T40
- helicini 204
- iliacus 217, 221
- inferior ossis pubis 196, 215, 234, 260, 266–267, 272–273, 317, 319
- infrapatellaris 348–349, 353, 365, T40
- intercostales anteriores 69, 247
- interganglionares 45
- interventricularis(-es) anterior (A. coronaria sinistra) 74, 79–80, 84, 86–87, 98, 127
- - posterior (A. coronaria dextra) 75, 80, 84–87, 120
- - septales (A. coronaria dextra) 86
- - - (A. coronaria sinistra) 86–87
- labiales posteriores 241
- lateralis (A. coronaria sinistra) 86
- - (N. spinalis, R. posterior) 41–43
- - (N. thoracicus) 37
- malleolares laterales 351, 372
- - mediales 351, 372
- mandibulae 12
- marginalis dexter 80, 84, 86–87
- - sinister 80, 84, 86–87
- mastoideus (A. occipitalis) 38
- medialis (N. spinalis, R. posterior) 41–43
- - (N. thoracicus) 37
- meningeus (Truncus n. spinalis) 41–43, 45
- muscularis(-es) (N. femoralis) 356–358, T40
- - (N. fibularis profundus) 374
- - (N. plantaris medialis) 377
- - (N. tibialis) 362–363, 369–370
- nodi atrioventriculares 86
- - sinuatrialis 84, 86
- obturatorius (A. epigastrica inferior) 225, 359
- occipitalis(-es) (A. auricularis posterior) 38
- - (A. occipitalis) 38

Ramus(-i)
- oesophageales (A. gastrica sinistra) 106, 131, 168
- - (A. thyroidea inferior) 106
- - (Aorta) 106, 115
- - (Ganglion thoracicum) 120
- - (N. laryngeus recurrens) 115, 121
- - (V. gastrica sinistra) 169, 181
- omentales (Aa. gastroomentales) 131, 172
- - (Vv. gastroomentales) 131, 160
- ossis ischii 234, 257, 261, 266–267, 273, 317, 319
- ovaricus 204
- perforans(-tes) (A. fibularis) 351, 368, 374
- - (A. thoracica interna) 69
- - (Arcus plantaris profundus) 378
- pericardiacus (N. phrenicus) 118
- perineales 240–241
- phrenicoabdominales (N. phrenicus) 63, 118
- posterior(-es) (A. media genus) 289
- - (A. obturatoria) 359
- - (A. renalis) 182
- - laterales (Nn. thoracici) 37
- - mediales (Nn. thoracici) 37
- - (N. auricularis magnus) 36
- - (N. cervicalis) 37–40
- - (N. obturatorius) 348, T40
- - (N. sacralis) 37
- - (N. spinalis) 36–37, 40–41, 45
- - (Nn. thoracici) 37–38, T20
- - (Truncus n. spinalis) 41–43, 45
- - (V. portae hepatis) 248
- - ventriculi sinistri 85–86
- posterolateralis dexter 86–87
- profundus (A. circumflexa femoris medialis) 359, 363–364
- - (A. glutea superior) 364
- - (A. plantaris medialis) 375
- - (A. transversa colli) 38–39
- - (N. plantaris lateralis) 377–378
- - (V. glutea superior) 364
- pubicus (A. epigastrica inferior) 225, 359
- - (A. obturatoria) 225
- radicularis anterior (A. spinalis anterior) 45
- saphenus (A. descendens genus) 351, 358
- scrotales anteriores 238
- - posteriores 227, 240
- sinister (A. hepatica propria) 146
- - (V. portae hepatis) 146, 154
- spinales (A. intercostalis posterior) 45
- - (A. sacralis lateralis) 226
- splenici 173
- subendocardiales 83
- superficialis (A. circumflexa femoris medialis) 357, 359, 363
- - (A. glutea superior) 363
- - (A. plantaris medialis) 375, 378
- - (A. transversa colli) 39
- - (N. plantaris lateralis) 376–378
- - (N. tibialis) T40
- superior ossis pubis 196, 234, 264, 266–267, 272–273, 317

Ramus(-i)
- tracheales (N. laryngeus recurrens) 120
- transversus 359
- tubarius (A. uterina) 204, 228
- vaginales (A. uterina) 204
Raphe
- perinei 232–233, 235
- scroti 194, 239
Recessus
- axillaris 122
- costodiaphragmaticus 54, 101, 103, 110, 118, 120–121, 160, 246–248, 250
- costomediastinalis 110, 113, 124, 127
- duodenalis inferior 164, 166–167
- - superior 164, 166–167
- ileocaecalis inferior 163–167
- - superior 163, 165–167
- inferior bursae omentalis 161
- intersigmoideus 164, 167
- phrenicomediastinalis 121, 243
- piriformis 108
- retrocaecalis 166
- splenicus 161, 167, 249
- subpopliteus 290, 331
- superior bursae omentalis 161, 167, 249
- suprapatellaris 292
- vertebromediastinalis 124, 126
Rectoscopy 211
Rectum 62, 138, 165–167, 177, 180, 185, 190, 205, 209–214, 220–221, 226, 228–229, 231, 237, 243, 256–257, 354
- AP-radiograph 141
- arteries 212
- colonoscopy 211
- innervation 214
- lateral radiograph 215
- rectoscopy 211
- veins 213
Regio(-nes)
- calcanea s. Calx
- cervicalis posterior 1, 38–40
- cruris 365
- - anterior 262, 367–368
- - posterior 262, 370–372
- deltoidea 1
- femoris anterior 262, 353, 356–358
- - posterior 262, 360–363
- genus anterior 262, 353
- - posterior 262
- glutealis 1, 262, 360–364
- infrascapularis 1
- inguinalis 262, 353–354
- lumbalis 1, 41
- - AP-radiograph 219
- occipitalis 38–39
- pedis 365
- perinealis of the female 241
- - of the male 240
- sacralis 1
- scapularis 1
- vertebralis 1
Ren 29, 35, 67, 152, 154–155, 160, 164, 166–167, 177, 179, 182–186, 188, 192, 200, 216–217, 245–247, 249–250, 255
- AP-radiograph 186, 216
- computed tomography 188
- development 188
- projection 183, 185
- ultrasound image 188
Rete
- articulare genus 351, 358, 368
- calcaneum 372, 377–378
- malleolare laterale 368, 374
- - mediale 374
- patellare 353, 356
- venosum dorsale pedis 365
Retinaculum(-a)
- musculorum extensorum inferius 306, 332, 334–337, 367–368, 373
- - - superius 306
- - fibularium inferius 304, 332, 334–336
- - - superius 304, 327, 329–332, 335, 370–371
- - flexorum (Pes) 304, 329–331, 335, 342, 370–371, 376–378
- - peroneorum inferius/superius s. Retinaculum musculorum fibularium inferius/superius
- patellae laterale 284–285, 306, 310–311
- - mediale 284–285, 323
Retroflexio (Uterus) 203
Retroperitoneal organs 152
Retroperitoneal space 179
Retroperitoneal structures, position of the 217–218
Retroversio (Uterus) 203
RETZIUS, Cave of 207, 252
Rima
- glottidis 114
- pudendi 235
Rims 5, 25
ROSENMÜLLER's node 355
Rugae vaginales 201, 205, 233

S

Sacral puncture 44
Sacroiliac joint s. Articulatio sacroiliaca
Sacrum, width of the 265
Salpinx s. Tuba uterina
SANTORINI's duct 152
Scalenus muscles T11
Scapula 1–3, 26–27, 48, 101, 108, 122–125
Scrotum 194, 197, 252
Segmentum(-a)
- anterius inferius (Ren) 187
- - laterale dextrum [VI] (Hepar) 147
- - - sinistrum [III] (Hepar) 147
- - mediale dextrum [V] (Hepar) 147
- - [S III] (Pulmo) 96
- - superius (Ren) 187
- apicale [S I] 96
- apicoposterius [S I-II] (Pulmo) 96
- basale anterius [S VIII] (Pulmo) 96
- - laterale [S IX] (Pulmo) 96
- - mediale [S VII] (Pulmo) 96
- - posterius [S X] (Pulmo) 96
- bronchopulmonalia 96–97
- cardiacum s. Segmentum basale mediale [S VII] (Pulmo)
- [I] (Hepar) 147
- inferius (Ren) 187
- laterale [S IV] (Pulmo) 96
- lingulare inferius [S V] (Pulmo) 96
- - superius [S IV] (Pulmo) 96
Segmentum(-a)
- mediale [S V] (Pulmo) 96
- - sinistrum [IV] 147
- posterius laterale dextrum [VII] (Hepar) 147
- - - sinistrum [II] (Hepar) 147
- - mediale dextrum [VIII] (Hepar) 147
- - (Ren) 187
- - [S II] (Pulmo) 96
- renalia 187
- superius (Ren) 187
- - [S VI] (Pulmo) 96
Septula testis 193–194, 252
Septum(-a)
- atrioventriculare 82
- femorale 355
- glandis 199
- interatriale 78–79, 119
- intermusculare cruris anterius 327, 332, 366, 381
- - - posterius 329, 366, 381
- - vastoadductorium 314, 357
- interventriculare 78–80, 83, 126
- longitudinalia 379
- penis 199, 243
- rectovaginale s. Fascia rectovaginalis
- scroti 61, 194, 239, 242
- vesicovaginale 207, 253
SHERMAN's vein 352
Sinus
- anales 210–211, 252
- aortae 79–80, 82, 127
- coronarius 75–76, 85, 99
- epididymidis 193–194
- lactiferus 64
- maxillaris 12
- obliquus pericardii 75, 89
- prostaticus 190
- renalis 183–184, 188, 250
- sagittalis superior 40
- tarsi 296, 305
- transversus 40
- - pericardii 75, 84, 89, 119, 126
- trunci pulmonalis 76, 82, 98
- venarum cavarum 85
Situs
- cordis 88
- retroperitonealis 179, 217–218
- viscerum 158–159, 162, 166, 172–173, 175
Sole of the foot, muscles 338–343, T53–T54
Spatium(-a)
- epidurale 24, 41, 43, 248, 252
- pararectale 205, 210
- paravesicale 205
- retropubicum 70, 205, 207, 252, 257
- retrorectale 205
- subarachnoideum 41, 43, 45, 119, 123, 252
- subdurale 41, 43
Spinal(-ae)
- iliaca anterior inferior 230, 264–267, 271, 273
- - - superior 3, 46–47, 52, 56–58, 71, 132, 140, 197, 207, 230, 255, 262, 264–269, 271, 273, 306, 315, 317, 322, 355, 360, 364
- - posterior inferior 266–267, 273
- - - superior 2, 31, 52, 183, 230, 254, 264, 266–267, 270, 273, 360
Spina(-ae)
- ischiadica 3, 212–213, 230, 256, 264, 266–267, 273, 280, 319
- scapulae 1–2, 26, 52, 103, 123
Spinal anaesthesia 44
Splen 140, 155–157, 159–161, 166–167, 170, 172, 180, 245, 247–249, 255
- AP-radiograph 171
- Hilum 156
Spondylolisthesis 9
Spondylolysis 9
Squama occipitalis 19
Sternoclavicular joint s. Articulatio sternoclavicularis
Sternum 5, 48, 50–52, 111, 119, 125, 127, 132, 207, 242
Stratum
- basale endometriale 202
- circulare 105, 129–130, 136, 139
- functionale endometriale 202
- longitudinale 105, 129–130, 136, 139, 210, 228
Stroma ovarii 201
Suboccipital muscles 31–34, 40, T20
Substantia
- compacta 277
- spongiosa 277, 281
Subtraction angiography, digital (DSA) 359
- arteries of femur 359
SUDECK's point 176, 212
Sulcus(-i)
- analis 262
- arteriae splenicae 153
- - subclaviae 49
- - vertebralis 6, 17
- calcanei 297–298
- coronarius 75–76, 85
- costae 49
- glutealis 262, 307, 319, 360
- interventricularis anterior 76, 80
- - posterior 76, 80, 85
- malleolaris 282–283, 301
- nervi spinalis 7
- obturatorius 266
- paracolici 162–164
- popliteus 281, 291, 293
- sinus sigmoidei 18
- supraacetabularis 266–267
- tali 297
- tendinis musculi fibularis longi 295, 297–298, 300, 302
- - - flexoris hallucis longi (Calcaneus) 296–298, 302
- - - - - longi (Talus) 297–298
- terminalis cordis 75, 85
- venae subclaviae 49
Sura 262
Sustentaculum tali 295–298, 300, 302, 305, 382
Sympathetic nervous system 214, 222–223
Symphysis
- manubriosternalis 50–51
- pubica 73, 190, 205, 207, 225, 230, 234–235, 244, 252–253, 256–257, 264, 268, 271–274, 317, 355, T39
- xiphosternalis 50
Synchondrosis costae 51
Syndesmosis tibiofibularis 263, 294, 304–305, 382, T39

Systema
- conducente cordis 82–83
- skeletale (Columna vertebralis) 2

T

Taenia
- libera 138–140, 158, 161–165
- mesocolica 139, 161
- omentalis 139, 158, 160–161, 163
Talus 295–299, 301, 304–305, 383
Tarsal joint
- s. Articulatio cuneonavicularis, Articulatio cuneocuboidea, Articulatio(-nes) intercuneiformes
- transverse s. Articulatio talocalcaneonavicularis, Articulatio calcaneocuboidea
Tarsal tunnel 377
Tarsometatarsal joints s. Articulatio(-nes) tarsometatarsales
Tarsus 263
Tela
- subcutanea 58, 74, 210, 232, 239
- submucosa 105, 130, 136–137, 139
- subserosa 130, 136, 139
Tendo calcaneus 300–301, 304, 307, 309, 328–333, 335, 370–372, 382–383
Testis 61, 67, 185, 192–194, 222, 242–243, 252
Thigh s. Femur
Thoracic muscles 52–55, 57–58, T13
Thoracic organs 118
Thoracic part of the vertebral column
- AP-radiograph 14
- lateral radiograph 14
Thoracic wall 58, 66, 69
- innervation, segmental 67
- muscles s. thoracic muscles
- skeleton projection 47
- surface anatomy 46
Thorax
- cross section 125
- frontal section 118, 121
- median section 119
Thymus 110–113, 119, 159
Tibia 262–263, 282–283, 285–287, 290–294, 300–301, 304–306, 327, 329–331, 366–367, 374, 381–383
Trabecula(-ae)
- carneae 79–80
- corporum cavernosorum 198
- septomarginalis 78, 80–81
- splenicae 156
Trachea 13, 89–91, 98–99, 101, 104, 106–107, 109, 111, 114–121, 123–124
- bifurcation 90
- cross-section 91
- projection 90
Tractus iliotibialis 279, 306–307, 312–313, 318–319, 322, 324, 327, 332, 361, 381
TREITZ, Muscle of 135, 152
Triad of GLISSON 145
Triangle of BOCHDALEK 63
Triangular space 37
Trigonum(-a)
- clavipectorale 58
- fibrosum dextrum 77, 84
- - sinistrum 77, 84
- lumbale 26–27, 37, 52, 364
Trigonum(-a)
- lumbocostale 63, T21
- sternocostale 63, T21
- vesicae 190–191, 195, 198
Trochanter
- major 256, 262, 268, 273–280, 319–321, 360, 364
- minor 273–280, 316, 321, 364
- tertius 275
Trochlea
- fibularis 296–297
- tali 295–298, 305
Truncus(-i)
- brachiocephalicus 74–75, 89, 98–99, 104–105, 107, 109, 111, 114–115, 118–121, 124
- coeliacus 62, 104, 106, 115, 152, 155, 168, 170–173, 178–179, 187, 214, 217–218, 220–221
- - variations 178
- costocervicalis 115
- (Embryo) 206
- gastrosplenicus 178
- hepatomesentericus 178
- intestinalis 218
- lumbalis 218–219
- lumbosacralis 68, 220–221, 225, 344
- nervi spinalis 41–43, 45
- pulmonalis 74–76, 79, 82, 84, 98, 101, 114, 120–121, 125–126
- sympathicus 41, 45, 63, 68, 83, 112–113, 115–117, 120–121, 124–125, 127, 220–223, 225, 245, 248, 250–251
- tibiofibularis 369, 372
- vagalis anterior 63, 117
- - posterior 63, 120–121
Trunk-arm muscles 26–27, T18
- deep layer 27
- superficial layer 26
Trunk-shoulder girdle muscles 26–27, T17
- deep layer 27
- superficial layer 26
Tuba
- uterina 167, 200–201, 203–204, 209, 223, 228, 259
- - AP-radiograph 203
Tuber
- calcanei 296–298, 300–302, 304–305, 328–330, 332, 340–341
- ischiadicum 3, 212–213, 215, 230, 232–235, 257–258, 265–268, 270, 273, 280, 317, 319–320, 360
- omentale 142, 161
Tubercle of VATER 135, 150, 152
Tuberculum(-a)
- adductorium 275, 277
- anterius (Vertebra cervicalis) 6–7, 35, 55
- costae 8, 20, 22, 49
- iliacum 264, 267
- intercondylare laterale 282, 291
- - mediale 282, 291
- intervenosum 78
- laterale (Talus) 296–298
- majus (Humerus) 47
- mediale (Talus) 296–298
- minus (Humerus) 47
- musculi scaleni anterioris 49
- obturatorium anterius 267
- - posterius 266–267
- posterius (Atlas) 34
- - (Vertebra cervicalis) 6–7, 12, 17, 31, 33, 35
Tuberculum(-a)
- pubicum 47, 230, 234–235, 264, 266–267, 271, 273, 355
- quadratum 275–276
Tuberositas
- glutea 275–276, 278, 319
- iliaca 264, 266
- musculi glutei maximi 267
- - serrati anterioris 49
- ossis cuboidei 295–296, 298, 302
- - metatarsi primi 295, 298
- - - quinti 295–296, 298, 302–303, 334
- - navicularis 295, 298, 302
- - sacri 10–11
- phalangis distalis (Pes) 295
- tibiae 262, 282, 285–287, 291, 294, 327, 332, 367
Tubulus
- distalis 184
- proximalis 184
Tunica
- adventitia 105
- - (Trachea) 91
- albuginea 193, 244
- - corporis spongiosi 199
- - corporum cavernosorum 197–199, 242, 252
- dartos 70, 194, 197, 226, 232, 239, 242, 252
- fibrosa 153
- mucosa 105, 121, 130, 136, 139, 150, 190, 201–202
- - (Gaster) 108
- - (Trachea) 91
- muscularis 105, 119, 129–130, 135–137, 139–140, 190–191, 201–202, 210, 226, 228, 230, 261
- serosa 130, 136–137, 143, 150, 153, 201–202, 226
- - (Colon) 139
- vaginalis testis 70, 193–194, 239, 242

U

Umbilical cord 208
Umbilicus 59, 72, 197
Uncovertebral clefts 24
Uncus corporis 7, 13, 24–25, 35
Upper abdomen
- vessels 172–173
- viscera 160
Urachus 70, 159, 185, 192, 200
Ureter 71, 152, 163–164, 166–167, 171, 179, 182–183, 185–187, 191–192, 196, 200–201, 205, 209, 216–218, 221–222, 226–229, 231, 236, 245, 251, 253–254, 258
- AP-radiograph 186
Urethra 190, 196, 198, 252–253
- feminina 200, 207, 230, 234, 237, 261
- masculina 190, 195–196, 198–199, 222, 234, 239, 244, 259–260
Uterus 140, 200–203, 205, 207, 209, 223, 228–229, 253, 354
- AP-radiograph 203
- mucosa, phase of proliferation 202
- position 203
- pregnant 206–207
- - ultrasound image 206
- ultrasound image 206
Utriculus prostaticus 190, 192, 198
Uvula vesicae 190, 198

V

Vagina(-ae) 200–205, 207, 223, 228, 230–231, 234, 237, 253, 257
- communis tendinum musculorum fibularium 304, 334–335
- musculi recti abdominis 52, 56–58, 69–70, 72, 249, 355
- plantaris tendinis musculi fibularis longi 341
- tendinis/tendinum digitorum pedis 335, 340–341
- - musculi/-orum extensoris digitorum longi 334–335
- - - - hallucis longi 334–335
- - - flexoris digitorum longi 304, 335
- - - - hallucis longi 335, 340
- - - tibialis anterioris 334–335
- - - - posterioris 304, 335
- - (Pes) 334–335
Valva(-ae)
- aortae 77, 80, 82, 85, 114, 119–120, 126–127
- atrioventricularis dextra 77–78, 80–82, 119–120, 127
- - sinistra 77, 79–82, 126–127
- cordis 77
- mitralis s. Valva atrioventricularis sinistra
- tricuspidalis s. Valva atrioventricularis dextra
- trunci pulmonalis 77, 82, 84
Valve of HEISTER 150
Valvula(-ae)
- anales 210
- foraminis ovalis 79
- fossae navicularis 198
- semilunaris anterior 77
- - - (Valva trunci pulmonalis) 84
- - dextra (Valva aortae) 77, 79, 82, 84, 119, 126
- - - (Valva trunci pulmonalis) 77, 84
- - posterior (Valva aortae) 77, 79–80, 82, 84, 119, 126
- - sinistra (Valva aortae) 77, 79–80, 82, 84, 119, 126
- - - (Valva trunci pulmonalis) 77, 82, 84
- sinus coronarii 78, 82, 84
- venae cavae inferioris 78, 82
Vas(-a)
- deferens s. Ductus deferens
- lymphaticum(-a) 197, 219
- - afferentia 219
- - AP-radiograph 219
- - centrale 136
- - efferentia 219
- - profunda 367
- - superficialia 366
- testiculares 71
VATER, Papilla of 135, 150, 152–153
Vein(s)
- deep perforating of HACH 352
- of BOYD 352
- of COCKETT 352
- of DODD 306, 352
- of HUNTER 352
- of MAY 352, 365
- of SHERMAN 352
Vena(-ae)
- anterior (V. pulmonalis dextra superior) 98, 120, 122
- apicalis 98
- apicoposterior 98
- appendicularis 169, 180

Vena(-ae)
- arcuata cruris posterior 365
- atriales sinistrae 75
- auricularis posterior 38–39
- axillaris 54, 113, 120, 122, 124
- azygos 63, 75, 99, 105–107, 112, 115–116, 121, 124–127, 181, 249
- basalis inferior 127
- basivertebralis 42, 119, 242
- brachiocephalica 75, 89, 98, 105–106, 110–113, 116, 118–119
- bronchialis 112
- bulbi penis 234
- - vestibuli 241
- cardiaca magna 74–75, 79–80, 84–85, 127
- - media 75, 84–85, 99, 119
- - parva 74, 85
- cava inferior 35, 63, 75–76, 78, 83, 85, 89, 99, 101, 105–107, 114, 116, 121, 142–143, 146–148, 152, 154–155, 167, 169, 172–173, 179–181, 185, 187–189, 210, 213, 217–218, 221, 243–244, 248–251
- - superior 74–78, 82, 84–85, 89, 98–99, 101, 105–107, 111–112, 114, 116, 118, 120, 124–126
- centralis 145, 189
- - (Glandula suprarenalis) 189
- cephalica 58, 66, 122
- cervicalis profunda 39
- - superficialis 66
- circumflexa femoris lateralis 261
- - - medialis 256, 260–261
- - humeri posterior 37
- - ilium profunda 179, 221, 227, 231
- - - superficialis 66, 353–355
- - scapulae 37
- colica dextra 152, 169, 175, 180
- - media 152, 161, 163–164, 169, 175, 177, 180
- - sinistra 167, 169, 175, 177, 180–181
- cordis 85
- - media s. Vena cardiaca media
- cremasterica 238
- cystica 149, 169, 172, 180–181
- digitales dorsales pedis 373
- dorsalis profunda clitoridis 231, 234
- - - penis 199, 227, 234, 238–239, 252
- - superficialis penis 199, 238, 252
- duodenales 169
- epigastrica inferior 60, 66, 69, 71–72, 158–159, 173, 175, 179, 181, 196, 217, 221, 227, 229, 231, 236, 253
- - superficialis 66, 181, 353–355
- - superior 66, 69–70, 72
- femoralis 62, 73, 218, 244, 256–257, 314, 353, 355–358, 380–381
- femoropoplitea 366
- fibularis 366
- gastrica(-ae) breves 131, 166, 169, 180
- - dextra 131, 169, 180
- - sinistra 106, 131, 152, 160, 169, 172, 179–181, 244, 249
- gastroduodenalis 152
- gastroomentalis dextra 131, 160, 167, 169, 172–173, 180
- - sinistra 106, 131, 160, 169, 172–173, 180
- glutea inferior 213, 256, 364, 380

Vena(-ae) glutea
- - superior 213, 254, 380
- hemiazygos 63, 106, 116, 119, 181, 245, 249
- - accessoria 106, 113, 116
- hepaticae 63, 89, 120–121, 143, 148, 167, 169, 179–181, 217–218
- - dextrae 146, 148, 243, 246, 248
- - intermediae 146, 148
- - projection 146
- - sinistrae 146, 148, 243–244
- - ultrasound image 148
- ileales 169, 175, 180, 251
- ileocolica 169, 180
- iliaca communis 167, 179, 181, 210, 213, 217–218, 221, 225, 227–228, 231, 242–243, 254
- - externa 71, 179, 196, 210, 213, 221, 226–229, 231, 236, 253–254, 317, 354, 356, 358
- - interna 181, 210, 213, 217, 226–228, 231, 254, 258, 354
- iliolumbalis 217
- intercostalis(-es) 70, 126–127
- - anteriores 247
- - posteriores 54, 73, 106, 112–113, 115–116
- interlobularis 145
- interventricularis anterior 74, 80, 84
- - posterior 75, 80, 84–85
- intervertebralis 42
- jejunales 152, 169, 175, 177, 180
- jugularis anterior 119
- - externa 38
- - interna 89, 105–106, 110, 118, 121–122
- lobi medii 98
- lumbales 41, 179, 217, 243, 245
- lumbalis ascendens 42, 115–116, 221
- marginalis coli 169
- - lateralis 365, 373
- - medialis 365, 373
- mesenterica inferior 152, 167, 169, 177, 179–181, 213
- - superior 152–153, 167, 169, 173, 175, 177, 179–181, 242–244, 251
- musculophrenica 69
- obliqua atrii sinistri 85
- obturatoria 71, 213, 227, 229, 231, 236, 256–257, 261, 317, 355
- occipitalis 37–39
- oesophageae 106, 169, 180
- ovarica 152, 167, 182, 187, 201, 209, 228–229, 253
- pancreaticae 169
- pancreaticoduodenalis(-es) 169, 180
- - superior posterior 169
- paraumbilicales 66, 72, 181
- perforantes 306, 352, 365, 373
- pericardiacophrenica 89, 111–113, 118
- perinealis 234
- phrenicae inferiores 63, 106, 181, 217
- plantaris lateralis 304
- - medialis 304
- poplitea 284, 329, 361–363, 366, 369–371
- portae hepatis 106, 131, 142–143, 146, 148, 152–154, 167, 169–170, 172–173, 179–181, 243–244, 246, 248–250
- - - projection 146

Vena(-ae) portae hepatis
- - - ultrasound image 148
- posterior (V. pulmonalis dextra superior) 98
- prepylorica 172
- profunda(-ae) brachii 37
- - femoris 261, 352, 358
- - interna 213, 227–228
- pudenda(-ae) externae 66, 197, 238, 353–355
- - interna 227, 231, 234, 236, 240–241, 256–258, 260–261, 363–364
- pulmonalis(-es) 112, 120
- - dextra(-ae) 75–76, 78–79, 85, 88–89, 95, 98, 100, 105, 107, 112, 121, 245
- - - inferior 99, 126–127
- - - superior 98–99, 120, 124, 126
- - sinistra(-ae) 75–76, 85, 89, 105, 107, 113, 119, 245
- - - inferior 79, 88, 95, 98–99, 126–127
- - - superior 79, 88, 95, 98–99, 107, 125
- rectalis(-es) inferiores 169, 181, 210, 213, 227–228
- - mediae 169, 210, 213, 227–229
- - superior 169, 177, 180–181, 210, 213, 227–229
- renalis 152, 154, 181–185, 187–188, 217, 221, 242–243, 245–246, 250
- sacralis lateralis 226
- - mediana 179, 210, 217, 221, 229
- saphena accessoria lateralis 353
- - - medialis 354
- - magna 66, 68, 218, 257, 284, 306, 352–356, 358, 360, 365–366, 373, 379, 381–382
- - parva 284, 352, 360–361, 363, 365–366, 369–370, 373, 381–382
- sigmoideae 169, 177, 179–181, 210, 213
- spinalis posterior 41
- splenica 152–156, 161, 167, 169, 173, 177, 179–181, 244–245, 248–250
- subclavia 89, 105–106, 111–112, 116, 118, 121, 123
- subcostalis 115–116
- subcutaneae abdominis 66
- suprarenalis 182, 187, 217
- suralis 370
- testicularis 71, 151, 179, 182, 187, 193, 217, 221, 238, 251
- thoracica interna 63, 66, 69, 88–89, 98, 110–111, 124
- - lateralis 66
- thoracoepigastricae 66
- thyroidea inferior 88–89, 106, 110–111, 116
- - superior 110
- tibiales anteriores 366–367
- - posteriores 366, 370
- transversa colli 38
- umbilicalis 70–71, 143, 159, 207
- uterina 228–229
- ventriculi dextri anteriores 84
- - sinistri posteriores 75, 85
- vertebralis 39, 75, 106
- vesicalis(-es) 70, 228
- - inferior 227

Vena(-ae) vesicalis(-es)
- - superior 227

Venter
- occipitalis 39
- posterior 31, 33–34
- superior 110

Ventriculus(-i)
- cordis dexter 74–76, 78, 80–82, 85, 98–99, 114, 120–121, 126–127, 244
- - sinister 74–76, 79–82, 85, 98, 101, 105, 120–121, 126–127

Versio (Uterus) 203

Vertebra(-ae)
- cervicales 3–5, 7, 12–13, 17–19, 35, 40, 48, 55, 103, 119
- - AP-radiograph 13
- - lateral radiograph 12
- coccygeae s. Os(-sa) coccygis
- development 5, 24
- epiphyses 5, 25
- lumbales 1, 3–5, 8–9, 15–16, 23–24, 32, 41, 44, 48, 62–63, 72, 154, 178, 216, 249–252, 254, 268–269, 271–273, 317
- - AP-radiograph 16
- - lateral radiograph 15
- ossification centres 5
- prominens 1–4, 7, 26, 34, 103, 119
- regional characteristics 5
- sacrales 2, 274
- structure characteristics 8
- thoracicae 3–5, 8–9, 20, 26, 48, 121–127, 186, 248–249
- - AP-radiograph 14
- - lateral radiograph 14

Vertebral column 3–4
- ligaments 20–21

Vesica
- biliaris 67, 140, 142, 144, 146–147, 149–151, 154, 160–161, 170, 172, 180, 250
- - AP-radiograph 151, 154
- - laparoscopy 149
- fellea s. Vesica biliaris
- urinaria 62, 67, 70–71, 140, 159, 165–167, 179, 185, 190–192, 196, 198, 200, 205, 207, 209, 217, 222, 226–229, 236–237, 242–243, 252–253, 256–257, 259–261, 354
- - puncture 190

Vestibulum
- bursae omentalis 161, 166
- vaginae 235

Villi intestinales 136

Viscera
- of the neonate 159
- upper abdomen 160–161

Vortex cordis 76

W

WINSLOW, Foramen of 160
WIRSUNG, Duct of 152
WOLFFian duct 192, 200

Z

ZENKER, Diverticulum of 109
Zona
- fasciculata 189
- orbicularis 256, 271, 279
- reticularis 189

Zygapophysis s. Processus articularis superior/inferior (Vertebra)